

SUPPLEMENT TO “A STRUCTURAL MODEL OF DENSE NETWORK FORMATION: APPENDICES A–B”
(*Econometrica*, Vol. 85, No. 3, May 2017, 825–850)

BY ANGELO MELE

APPENDIX A: MODEL EQUILIBRIUM: PROOFS

PROOF OF PROPOSITION 1: The potential is a function Q from the space of actions to the real line such that $Q(g_{ij}, g_{-ij}, X) - Q(g'_{ij}, g_{-ij}, X) = U_i(g_{ij}, g_{-ij}, X) - U_i(g'_{ij}, g_{-ij}, X)$, for any ij .⁴⁵ A simple computation shows that, for any ij ,

$$\begin{aligned} & Q(g_{ij} = 1, g_{-ij}, X) - Q(g_{ij} = 0, g_{-ij}, X) \\ &= u_{ij} + g_{ji}m_{ij} + \sum_{\substack{k=1 \\ k \neq i, j}}^n g_{jk}v_{ik} + \sum_{\substack{k=1 \\ k \neq i, j}}^n g_{ki}v_{kj} \\ &= U_i(g_{ij} = 1, g_{-ij}, X) - U_i(g_{ij} = 0, g_{-ij}, X); \end{aligned}$$

therefore, Q is the potential of the network formation game. *Q.E.D.*

PROOF OF COROLLARY 1: The proof consists of showing that $Q(g, X)$ can be written in the form $\theta' \mathbf{t}(g, X)$. Consider the first part of the potential:

$$\begin{aligned} \sum_i \sum_j g_{ij} u_{ij} &= \sum_i \sum_j g_{ij} \sum_{p=1}^P \theta_{up} H_{up}(X_i, X_j) \\ &= \sum_{p=1}^P \theta_{up} \sum_i \sum_j g_{ij} H_{up}(X_i, X_j) \\ &\equiv \sum_{p=1}^P \theta_{up} t_{up}(g, X) = \theta'_u \mathbf{t}_u(g, X), \end{aligned}$$

where $t_{up}(g, X) \equiv \sum_i \sum_j g_{ij} H_{up}(X_i, X_j)$, $\theta_u = (\theta_{u1}, \dots, \theta_{uP})'$, and $\mathbf{t}_u(g, X) = (t_{u1}(g, X), \dots, t_{uP}(g, X))'$. Analogously define $\theta_m = (\theta_{m1}, \theta_{m2}, \dots, \theta_{mL})'$ and $\mathbf{t}_m(g, X) = (t_{m1}(g, X), t_{m2}(g, X), \dots, t_{mL}(g, X))'$ and $\theta_v = (\theta_{v1}, \theta_{v2}, \dots, \theta_{vS})'$ and $\mathbf{t}_v(g, X) = (t_{v1}(g, X), t_{v2}(g, X), \dots, t_{vS}(g, X))'$. It follows that

$$\begin{aligned} \sum_i \sum_{j>i} g_{ij} g_{ji} m_{ij} &= \theta'_m \mathbf{t}_m(g, X), \\ \sum_i \sum_j g_{ij} \sum_{k \neq i, j} g_{jk} v_{ij} &= \theta'_v \mathbf{t}_v(g, X). \end{aligned}$$

⁴⁵For more details and definitions, see Monderer and Shapley (1996).

Therefore, $Q(g, X)$ can be written in the form $\boldsymbol{\theta}'\mathbf{t}(g, X)$, where $\boldsymbol{\theta} = (\boldsymbol{\theta}_u, \boldsymbol{\theta}_m, \boldsymbol{\theta}_v)'$ and $\mathbf{t}(g, X) = [\mathbf{t}_u(g, X), \mathbf{t}_m(g, X), \mathbf{t}_v(g, X)]'$,

$$\begin{aligned} Q(g, X) &= \boldsymbol{\theta}'_u \mathbf{t}_u(g, X) + \boldsymbol{\theta}'_m \mathbf{t}_m(g, X) + \boldsymbol{\theta}'_v \mathbf{t}_v(g, X) \\ &= \boldsymbol{\theta}'\mathbf{t}(g, X) \end{aligned}$$

and the result follows. Q.E.D.

Model Without Preference Shocks: Nash Networks

Let us consider a *special case* of the model, in which there are no preference shocks. Let $\mathcal{N}(g)$ be the set of networks that differ from g by only one element of the matrix, that is,

$$\mathcal{N}(g) \equiv \{g' : g' = (g'_{ij}, g_{-ij}), \text{ for all } g'_{ij} \neq g_{ij}, \text{ for all } i, j \in \mathcal{I}\}. \quad (19)$$

A Nash network is defined as a network in which any player has no profitable deviations from his current linking strategy, when randomly selected from the population. The following results characterize the set of the pure-strategy Nash equilibria and the long-run behavior of the model with no shocks.

PROPOSITION 2—Model Without Shocks: Equilibria and Long Run: *Consider the model without idiosyncratic preference shocks. Under Assumptions 1 and 2:*

1. *There exists at least one pure-strategy Nash equilibrium network.*
2. *The set $\mathcal{NE}(\mathcal{G}, X, U)$ of all pure-strategy Nash equilibria of the network formation game is completely characterized by the local maxima of the potential function:*

$$\mathcal{NE}(\mathcal{G}, X, U) = \left\{ g^* : g^* = \arg \max_{g \in \mathcal{N}(g^*)} Q(g, X) \right\}. \quad (20)$$

3. *Any pure-strategy Nash equilibrium is an absorbing state.*
4. *As $t \rightarrow \infty$, the network converges to one of the Nash networks with probability 1.*

PROOF: (1) The existence of Nash equilibria follows directly from the fact that the network formation game is a potential game with finite strategy space (see [Monderer and Shapley \(1996\)](#) for details).

(2) The set of Nash equilibria is defined as the set of g^* such that, for every i and for every $g_{ij} \neq g^*_{ij}$,

$$U_i(g^*_{ij}, g^*_{-ij}, X) \geq U_i(g_{ij}, g^*_{-ij}, X).$$

Therefore, since Q is a potential function, for every $g_{ij} \neq g^*_{ij}$,

$$Q(g^*_{ij}, g^*_{-ij}, X) \geq Q(g_{ij}, g^*_{-ij}, X).$$

Therefore, g^* is a maximizer of Q . The converse is easily checked by the same reasoning.

(3) Suppose $g^t = g^*$. Since this is a Nash equilibrium, no player will be willing to change her linking decision when her turn to play comes. Therefore, once the chain reaches a Nash equilibrium, it cannot escape from that state.

(4) The probability that the potential will increase from t to $t + 1$ is

$$\begin{aligned}
 & \Pr[Q(g^{t+1}, X) \geq Q(g^t, X)] \\
 &= \sum_i \sum_j \Pr(m^{t+1} = ij) \underbrace{\Pr[U_i(g_{ij}^{t+1}, g_{-ij}^t, X) \geq U_i(g_{ij}^t, g_{-ij}^t, X) | m^{t+1} = ij]}_{=1 \text{ because agents play Best Response, conditioning on } m^{t+1}} \\
 &= \sum_i \sum_j \rho_{ij} = 1.
 \end{aligned}$$

By part (3) of the proposition, a Nash network is an absorbing state of the chain. Therefore, any probability distribution that puts probability 1 on a Nash network is a stationary distribution. For any initial network, the chain will converge to one of the stationary distributions. It follows that, in the long run, the model will be in a Nash network, that is, for any $g^0 \in \mathcal{G}$,

$$\lim_{t \rightarrow \infty} \Pr[g^t \in NE | g^0] = 1. \quad \text{Q.E.D.}$$

PROOF OF THEOREM 1: 1. The sequence of networks $[g^0, g^1, \dots]$ generated by the network formation game is a Markov chain. Inspection of the transition probability proves that the chain is irreducible and aperiodic; therefore, it is ergodic. The existence of a unique stationary distribution then follows from the ergodic theorem (see Gelman, Roberts, and Gilks (1996) for details).

2. A sufficient condition for stationarity is the *detailed balance* condition. In our case, this requires $P_{gg'} \pi_g = P_{g'g} \pi_{g'}$, where $P_{gg'} = \Pr(g^{t+1} = g' | g^t = g)$ and $\pi_g = \pi(g^t = g)$. The transition from g to g' is possible if these networks differ by only one element g_{ij} . Otherwise, the transition probability is zero and the detailed balance condition is satisfied. Let us consider the nonzero probability transitions, with $g = (1, g_{-ij})$ and $g' = (0, g_{-ij})$. Define $\Delta Q \equiv Q(1, g_{-ij}, X) - Q(0, g_{-ij}, X)$:

$$\begin{aligned}
 P_{gg'} \pi_g &= \Pr(m^t = ij) \Pr(g_{ij} = 0 | g_{-ij}) \frac{\exp[Q(1, g_{-ij}, X)]}{\sum_{\omega \in \mathcal{G}} \exp[Q(\omega, X)]} \\
 &= \rho(g_{-ij}, X_i, X_j) \times \frac{1}{1 + \exp[\Delta Q]} \\
 &\quad \times \frac{\exp[Q(1, g_{-ij}, X) + Q(0, g_{-ij}, X) - Q(0, g_{-ij}, X)]}{\sum_{\omega \in \mathcal{G}} \exp[Q(\omega, X)]} \\
 &= \rho(g_{-ij}, X_i, X_j) \times \frac{1}{1 + \exp[\Delta Q]} \\
 &\quad \times \frac{\exp[Q(1, g_{-ij}, X) - Q(0, g_{-ij}, X)] \exp[Q(0, g_{-ij}, X)]}{\sum_{\omega \in \mathcal{G}} \exp[Q(\omega, X)]} \\
 &= \rho(g_{-ij}, X_i, X_j) \frac{\exp[\Delta Q]}{1 + \exp[\Delta Q]} \frac{\exp[Q(0, g_{-ij}, X)]}{\sum_{\omega \in \mathcal{G}} \exp[Q(\omega, X)]}
 \end{aligned}$$

$$\begin{aligned}
&= \Pr(m' = ij) \Pr(g_{ij} = 1 | g_{-ij}) \frac{\exp[Q(0, g_{-ij}, X)]}{\sum_{\omega \in \mathcal{G}} \exp[Q(\omega, X)]} \\
&= P_{g'g} \pi_{g'}.
\end{aligned}$$

So the distribution (5) satisfies the detailed balance condition. Therefore, it is a stationary distribution for the network formation model. From part (1) of the proposition, we know that the process is ergodic and it has a unique stationary distribution. Therefore, $\pi(g, X)$ is also the unique stationary distribution. *Q.E.D.*

APPENDIX B: COMPUTATIONAL DETAILS

B.1. Network Simulation

The algorithm used to simulate the network (Algorithm 1) produces samples from the stationary equilibrium of the model.

1. The network simulation algorithm satisfies the detailed balance condition for the stationary distribution (5). Indeed, for any given θ ,

$$\begin{aligned}
\Pr(g'|g, X, \theta) \pi(g, X, \theta) &= q_g(g'|g) \min \left\{ 1, \frac{\exp[Q(g', X, \theta)] q_g(g|g')}{\exp[Q(g, X, \theta)] q_g(g'|g)} \right\} \frac{\exp[Q(g, X, \theta)]}{c(\mathcal{G}, X, \theta)} \\
&= \min \left\{ q_g(g'|g) \frac{\exp[Q(g, X, \theta)]}{c(\mathcal{G}, X, \theta)}, \frac{\exp[Q(g', X, \theta)]}{c(\mathcal{G}, X, \theta)} q_g(g|g') \right\} \\
&= q_g(g|g') \min \left\{ \frac{q_g(g'|g) \exp[Q(g, X, \theta)]}{q_g(g|g') c(\mathcal{G}, X, \theta)}, \frac{\exp[Q(g', X, \theta)]}{c(\mathcal{G}, X, \theta)} \right\} \\
&= q_g(g|g') \min \left\{ \frac{q_g(g'|g) \exp[Q(g, X, \theta)]}{q_g(g|g') \exp[Q(g', X, \theta)]}, 1 \right\} \frac{\exp[Q(g', X, \theta)]}{c(\mathcal{G}, X, \theta)} \\
&= \Pr(g|g', X, \theta) \pi(g', X, \theta).
\end{aligned}$$

This concludes the proof.

2. The algorithm generates a Markov chain of networks with finite state space. The chain is irreducible and aperiodic and therefore it is uniformly ergodic (see Theorem 4.9, p. 52 in [Levin, Peres, and Wilmer \(2008\)](#)).

3. The bound to the convergence rate used in the text was derived by [Diaconis and Stroock \(1991\)](#), for reversible finite chains.

The algorithm has a very useful property that can be exploited in the posterior simulation to reduce the computational burden. Adapting the suggestion in [Liang \(2010\)](#), define $\mathcal{P}_{\theta'}^{(R)}(g'|g)$ as the transition probability of a Markov chain that generates g' with R Metropolis–Hastings updates of the network simulation algorithm, starting at the observed network g and using the proposed parameter θ' . Then,

$$\mathcal{P}_{\theta'}^{(R)}(g'|g) = \mathcal{P}_{\theta'}(g^1|g) \mathcal{P}_{\theta'}(g^2|g^1) \cdots \mathcal{P}_{\theta'}(g'|g^{R-1}), \quad (21)$$

where $\mathcal{P}_{\theta'}(g^j|g^i) = q_g(g^j|g^i) \alpha_{mh}(g^i, g^j)$ is the transition probability of the network simulation algorithm above. Since the Metropolis–Hastings algorithm satisfies the detailed balance condition, we can prove the following lemma.

LEMMA 1: Simulate a network g' from the stationary distribution $\pi(\cdot, X, \theta')$ using a Metropolis–Hastings algorithm starting at the network g observed in the data. Then

$$\frac{\mathcal{P}_{\theta'}^{(R)}(g|g')}{\mathcal{P}_{\theta'}^{(R)}(g'|g)} = \frac{\exp[Q(g, X, \theta')]}{\exp[Q(g', X, \theta')]} \quad (22)$$

for all R , $g, g' \in \mathcal{G}$ and for any $\theta' \in \Theta$.

PROOF: Let $\mathcal{P}_{\theta'}^{(R)}(g'|g)$ be defined as in (21). This is the transition probability of the chain that generates g' with R Metropolis–Hastings updates, starting at the observed network g and using the proposed parameter θ' . Notice that the Metropolis–Hastings algorithm satisfies the detailed balance for $\pi(g, X, \theta')$; therefore, we have

$$\begin{aligned} \mathcal{P}_{\theta'}^{(R)}(g|g')\pi(g', X, \theta') &= \mathcal{P}_{\theta'}(g_{R-1}|g')\mathcal{P}_{\theta'}(g_{R-2}|g_{R-1}) \cdots \mathcal{P}_{\theta'}(g|g_1)\pi(g', X, \theta') \\ &= \mathcal{P}_{\theta'}(g_1|g)\mathcal{P}_{\theta'}(g_2|g_1) \cdots \mathcal{P}_{\theta'}(g'|g_{R-1})\pi(g, X, \theta') \\ &= \mathcal{P}_{\theta'}^{(R)}(g'|g)\pi(g, X, \theta'). \end{aligned}$$

It follows that

$$\begin{aligned} \frac{\mathcal{P}_{\theta'}^{(R)}(g|g')}{\mathcal{P}_{\theta'}^{(R)}(g'|g)} &= \frac{\pi(g, X, \theta')}{\pi(g', X, \theta')} \\ &= \frac{\exp[Q(g, X, \theta')]}{\exp[Q(g', X, \theta')]} \frac{c(\mathcal{G}, X, \theta')}{c(\mathcal{G}, X, \theta')} \\ &= \frac{\exp[Q(g, X, \theta')]}{\exp[Q(g', X, \theta')]} \quad \text{Q.E.D.} \end{aligned}$$

One should notice that as long as the algorithm is started from the network g observed in the data (which is assumed to be a draw from the stationary equilibrium of the model), the equality in (22) is satisfied for any R .

The approximate exchange algorithm presented in this paper removes the requirement of exact sampling by exploiting the property of the stationary equilibrium characterization, described in Lemma 1.

Local Simulations and Large Steps

The theoretical results using graph limits and large deviations suggest that the local sampler has systematic convergence problems, even when in principle the simulation is trivial because links are asymptotically independent ($\beta > 0$).

To attenuate these convergence issues, we propose a modification that allows the sampler to make *larger steps*, in particular, steps that are not $o(n)$. The local chain selects a link g_{ij} with probability $1/(n(n-1))$, proposing to swap the value to $1 - g_{ij}$. We add the following large steps. First, with probability p_r , the sampler selects a player i at random (with probability $1/n$) and proposes to swap all his links, that is, $g_{ij} = 1 - g_{ij}$ for each $j = 1, \dots, n$. Second, with probability p_c , the sampler selects a player i at random (with probability $1/n$) and proposes to swap all the links pointing at i , that is, $g_{ji} = 1 - g_{ji}$ for each $j = 1, \dots, n$. Third, with probability p_f , the sampler selects uniformly at random

$[\lambda n]$ links, where $\lambda \in (0, 1)$, and proposes to swap all of them. Notice that this step size is a function of n , and in particular is not $o(n)$. The crucial ingredient is to make the length of the step a function of n . The parameter λ is under control of the researcher: higher values allow larger steps and increase the computational cost of sampling. Last, with probability p_{inv} , the sampler proposes to invert the adjacency matrix. The goal of this large step is to provide a way to jump across modes of the stationary distribution, when it is bimodal.⁴⁶

Using this sampler, we reproduce the simulation in Figure 1. We know that the local chain can get trapped in local maxima of the variational problem. If we simulate model (10) with parameters $(\alpha, \beta) = (-3, 3)$, we obtain Figure 8(A). While Theorem 2 states that the simulations should converge to the sparse network density $\mu_1 \approx 0.07$, we observe that the local sampler converges to a dense network with $\mu_2 \approx 0.93$, if started at dense networks. In other words, when started at a dense network (say the full network), the sampler gets trapped in a local maximum of the variational problem, with density $\mu_2 \approx 0.93$. Figure 8(B) shows that our modified sampler does not have this problem, and also the chains started at dense networks converge to the correct (sparse) network density. This simple modification gets rid of the exponentially slow convergence of the local algorithm. More generally, these larger steps allow the sampler to escape local maxima of the potential function.

In general, this modification should help the sampler when the likelihood has multiple modes. However, the improvement comes with the increased cost of sampling and additional computational time. In some models, the cost may be substantial. For example, it is intuitive that in regions of the parameter space (say, $\beta < 0$) where the likelihood is unimodal, the gains from this modified sampler are minimal.




FIGURE 8.—Local sampler versus modified sampler. Comparison of network samplers for model (10), with parameters $(\alpha, \beta) = (-3, 3)$. Panel (A) shows the simulation using the local-chain sampler, which converges to two different link densities ($\mu_1 \approx 0.07$ and $\mu_2 \approx 0.93$). However, we know from Theorem 2 that the correct simulation should converge to the sparse network density. So the local chain fails to sample correctly if we start it at a dense network, because it gets trapped at a local maximum of the stationary distribution. Panel (B) shows the simulation using the modified algorithm. We use $p_r = p_f = p_{\text{inv}} = 0.01$. The simulations converge to the correct link density for any starting value; therefore, our modified algorithm provides a better sampler for the model.

⁴⁶We have seen that this is the case in the homogeneous player case, for many parameter values.

B.2. Posterior Simulation

In this section, I provide the technical details for the algorithm proposed in the empirical part of the paper. The first set of results show that the exchange algorithm generates (approximate) samples from the posterior distribution (7).

The original exchange algorithm developed in Murray, Ghahramani, and MacKay (2006) is slightly different from the one used here. The main modification is in Step 2: the original algorithm requires an *exact* sample from the stationary equilibrium of the model.

ALGORITHM 3—Exact Exchange Algorithm: Start at current parameter $\theta_t = \theta$ and network data g .

1. Propose a new parameter vector θ'

$$\theta' \sim q_\theta(\cdot|\theta). \quad (23)$$

2. Draw an exact sample network g' from the likelihood

$$g' \sim \pi(\cdot|X, \theta'). \quad (24)$$

3. Compute the acceptance ratio

$$\begin{aligned} \alpha_{ex}(\theta, \theta', g', g) &= \min \left\{ 1, \frac{\exp[Q(g', X, \theta)]}{\exp[Q(g, X, \theta)]} \frac{p(\theta')}{p(\theta)} \frac{q_\theta(\theta|\theta')}{q_\theta(\theta'|\theta)} \frac{\exp[Q(g, X, \theta')]}{\exp[Q(g', X, \theta')]} \frac{c(\theta)c(\theta')}{c(\theta)c(\theta')} \right\} \quad (25) \\ &= \min \left\{ 1, \frac{\exp[Q(g', X, \theta)]}{\exp[Q(g, X, \theta)]} \frac{p(\theta')}{p(\theta)} \frac{q_\theta(\theta|\theta')}{q_\theta(\theta'|\theta)} \frac{\exp[Q(g, X, \theta')]}{\exp[Q(g', X, \theta')]} \right\}. \end{aligned}$$

4. Update the parameter according to

$$\theta_{t+1} = \begin{cases} \theta', & \text{with prob. } \alpha_{ex}(\theta, \theta', g', g), \\ \theta, & \text{with prob. } 1 - \alpha_{ex}(\theta, \theta', g', g). \end{cases} \quad (26)$$

The difference between this algorithm and the approximate one is in step 2. The exact and approximate algorithms use the same acceptance ratio $\alpha_{ex}(\theta, \theta', g', g)$, a consequence of Lemma 1. Indeed, the acceptance ratio for the approximate algorithm is

$$\tilde{\alpha}_{ex}(\theta, \theta', g', g) = \min \left\{ 1, \frac{\exp[Q(g', X, \theta)]}{\exp[Q(g, X, \theta)]} \frac{p(\theta')}{p(\theta)} \frac{q_\theta(\theta|\theta')}{q_\theta(\theta'|\theta)} \frac{\mathcal{P}_{\theta'}^{(R)}(g|g')}{\mathcal{P}_{\theta'}^{(R)}(g'|g)} \right\} \quad (27)$$

$$= \min \left\{ 1, \frac{\exp[Q(g', X, \theta)]}{\exp[Q(g, X, \theta)]} \frac{p(\theta')}{p(\theta)} \frac{q_\theta(\theta|\theta')}{q_\theta(\theta'|\theta)} \frac{\exp[Q(g, X, \theta')]}{\exp[Q(g', X, \theta')]} \right\} \quad (28)$$

$$= \alpha_{ex}(\theta, \theta', g', g). \quad (29)$$

This result implies that to prove the convergence of the approximate algorithm to the exact algorithm, there is no need to prove convergence of $\tilde{\alpha}_{ex}(\theta, \theta', g', g)$ to $\alpha_{ex}(\theta, \theta', g', g)$. The convergence of step 2 of the algorithm is sufficient.

B.2.1. Preliminary Lemmas for Theorem 6

The convergence of the approximate exchange algorithm to the correct posterior distribution is proven in four steps. First, we prove that the exact exchange algorithm converges to the correct posterior (Lemma 2). Second, we prove that the approximate algorithm has a stationary distribution and it is ergodic (Lemma 3, similar to Liang (2010)). Third, we prove that the transition kernels of the approximate and exact algorithms are arbitrarily close for a large enough number of network simulations (Lemma 4). Fourth, we combine previous results to prove that the approximate algorithm converges to the correct posterior. A similar proof strategy is contained in Liang, Liu, and Carroll (2010) and Andrieu and Roberts (2009).

Let $Q(d\vartheta|\theta) = q_\theta(\vartheta|\theta)\nu(d\vartheta)$. The transition kernel of the exact exchange algorithm can be written as

$$\begin{aligned} P(\theta, d\vartheta) &= \left[\sum_{g' \in \mathcal{G}} \pi(g', \vartheta) \alpha_{ex}(\theta, \vartheta, g', g) \right] Q(\theta, d\vartheta) \\ &\quad + \delta_\theta(d\vartheta) \left\{ 1 - \int_{\Theta} \left[\sum_{g' \in \mathcal{G}} \pi(g', \vartheta) \alpha_{ex}(\theta, \vartheta, g', g) \right] Q(\theta, d\vartheta) \right\} \end{aligned}$$

and the transition kernel of the approximate exchange algorithm can be written as

$$\begin{aligned} \tilde{P}_R(\theta, d\vartheta) &= \left[\sum_{g' \in \mathcal{G}} \mathcal{P}_\vartheta^{(R)}(g'|g) \alpha_{ex}(\theta, \vartheta, g', g) \right] Q(\theta, d\vartheta) \\ &\quad + \delta_\theta(d\vartheta) \left\{ 1 - \int_{\Theta} \left[\sum_{g' \in \mathcal{G}} \mathcal{P}_\vartheta^{(R)}(g'|g) \alpha_{ex}(\theta, \vartheta, g', g) \right] Q(\theta, d\vartheta) \right\}. \end{aligned}$$

Let $\eta(\theta)$ be the average rejection probability for the approximate algorithm, that is,

$$\eta(\theta) := 1 - \int_{\Theta} \left[\sum_{g' \in \mathcal{G}} \mathcal{P}_\vartheta^{(R)}(g'|g) \alpha_{ex}(\theta, \vartheta, g', g) \right] Q(\theta, d\vartheta). \quad (30)$$

The next lemma proves that the transition kernel satisfies the detailed balance condition for the posterior distribution. For any pair of parameters $(\theta, \vartheta) \in \Theta$, we have

$$P[\theta, \vartheta|g, X]p(\theta|g, X) = \Pr[\theta|\vartheta, g, X]p(\vartheta|g, X). \quad (31)$$

The detailed balance condition is sufficient to prove that posterior (7) is the stationary distribution of the Markov chain generated by the algorithm (for details, see Robert and Casella (2005) or Gelman et al. (2003)).

LEMMA 2: *The exchange algorithm produces a Markov chain with invariant distribution (7).*

PROOF: Define $\mathcal{Z} \equiv \int_{\Theta} \pi(g|X, \theta)p(\theta) d\theta$. In the algorithm, the probability $\Pr[\vartheta|\theta, g, X]$ of transition to θ_j , given the current parameter θ and the observed data (g, X) , can be computed as

$$\Pr[\vartheta|\theta, g, X] = q_\theta(\vartheta|\theta) \frac{\exp[Q(g', X, \vartheta)]}{c(\mathcal{G}, X, \vartheta)} \alpha_{ex}(\theta, \vartheta, g', g). \quad (32)$$

This is the probability $q_\theta(\vartheta|\theta)$ of proposing ϑ times the probability of generating the new network g' from the model's stationary distribution, $\frac{\exp[Q(g', X, \vartheta)]}{c(\mathcal{G}, X, \vartheta)}$, and accepting the proposed parameter $\alpha_{ex}(\theta, \vartheta, g', g)$. Therefore, the left-hand side of (31) can be written as

$$\begin{aligned}
 \Pr[\vartheta|\theta, g, X]p(\theta|g, X) &= q_\theta(\vartheta|\theta) \frac{\exp[Q(g', X, \vartheta)]}{c(\mathcal{G}, X, \vartheta)} \alpha_{ex}(\theta, \vartheta, g', g) p(\theta|g, X) \\
 &= q_\theta(\vartheta|\theta) \frac{\exp[Q(g', X, \vartheta)]}{c(\mathcal{G}, X, \vartheta)} \alpha_{ex}(\theta, \vartheta, g', g) \frac{\exp[Q(g, X, \theta)]}{c(\mathcal{G}, X, \theta)} p(\theta) \\
 &= q_\theta(\vartheta|\theta) \frac{\exp[Q(g', X, \vartheta)]}{c(\mathcal{G}, X, \vartheta)} \\
 &\quad \times \min \left\{ 1, \frac{\exp[Q(g', X, \theta)]}{\exp[Q(g, X, \theta)]} \frac{p(\vartheta)}{p(\theta)} \frac{q_\theta(\theta|\vartheta)}{q_\theta(\vartheta|\theta)} \frac{\exp[Q(g, X, \vartheta)]}{\exp[Q(g', X, \vartheta)]} \right\} \\
 &\quad \times \frac{\exp[Q(g, X, \theta)]}{c(\mathcal{G}, X, \theta)} p(\theta) \\
 &\quad \times \frac{1}{\mathcal{Z}} \\
 &= \min \left\{ q_\theta(\vartheta|\theta) \frac{\exp[Q(g', X, \vartheta)]}{c(\mathcal{G}, X, \vartheta)} \frac{\exp[Q(g, X, \theta)]}{c(\mathcal{G}, X, \theta)} \frac{p(\theta)}{\mathcal{Z}}, \right. \\
 &\quad \left. q_\theta(\theta|\vartheta) \frac{\exp[Q(g', X, \theta)]}{c(\mathcal{G}, X, \theta)} \frac{\exp[Q(g, X, \vartheta)]}{c(\mathcal{G}, X, \vartheta)} \frac{p(\vartheta)}{\mathcal{Z}} \right\} \\
 &= q_\theta(\theta|\vartheta) \frac{\exp[Q(g', X, \theta)]}{c(\mathcal{G}, X, \theta)} \frac{\exp[Q(g, X, \vartheta)]}{c(\mathcal{G}, X, \vartheta)} \frac{p(\vartheta)}{\mathcal{Z}} \\
 &\quad \times \min \left\{ 1, \frac{\exp[Q(g', X, \vartheta)]}{\exp[Q(g, X, \vartheta)]} \frac{p(\theta)}{p(\vartheta)} \frac{q_\theta(\vartheta|\theta)}{q_\theta(\theta|\vartheta)} \frac{\exp[Q(g, X, \theta)]}{\exp[Q(g', X, \theta)]} \right\} \\
 &= q_\theta(\theta|\vartheta) \frac{\exp[Q(g', X, \theta)]}{c(\mathcal{G}, X, \theta)} \alpha(\vartheta, \theta, g', g) \frac{\exp[Q(g, X, \vartheta)]}{c(\mathcal{G}, X, \vartheta)} \frac{p(\vartheta)}{\mathcal{Z}} \\
 &= q_\theta(\theta|\vartheta) \frac{\exp[Q(g', X, \theta)]}{c(\mathcal{G}, X, \theta)} \alpha(\vartheta, \theta, g', g) p(\vartheta|g, X) \\
 &= \Pr[\theta|\vartheta, g, X] p(\vartheta|g, X).
 \end{aligned}$$

The latter step proves the detailed balance for a generic network g' . Since the condition is satisfied for any network g' , detailed balance follows from summing over all possible networks. *Q.E.D.*

LEMMA 3—The Approximate Algorithm is Ergodic: *Assume the exact exchange algorithm is ergodic and that, for any $\vartheta \in \Theta$,*

$$\frac{\mathcal{P}_\vartheta^{(R)}(g'|g)}{\pi(g', \vartheta)} > 0 \quad \text{for any } g' \in \mathcal{G}. \quad (33)$$

Then, for any $R \in \mathbb{N}$ such that, for any $\theta \in \Theta$, $\rho(\theta) > 0$, the transition kernel of the approximate algorithm \tilde{P}_R is also irreducible and aperiodic, and there exists a stationary distribution $\tilde{p}(\theta)$ such that

$$\lim_{s \rightarrow \infty} \|\tilde{P}_R^{(s)}(\theta_0, \cdot) - \tilde{p}(\theta)\|_{TV} = 0. \quad (34)$$

PROOF: The exact algorithm with transition kernel P is an irreducible and aperiodic Markov chain. To prove that the approximate algorithm with transition kernel \tilde{P}_R defines an ergodic Markov chain, it is sufficient to prove that the set of accessible states of P are also included in those of \tilde{P}_R . The proof proceeds by induction.

Formally, we need to show that for any $s \in \mathbb{N}$, $\theta \in \Theta$ and $A \in \mathcal{B}(\Theta)$ such that $P^{(s)}(\theta, A) > 0$, implies $\tilde{P}_R^{(s)}(\theta, A) > 0$.

Notice that for any $\theta \in \Theta$ and $A \in \mathcal{B}(\Theta)$,

$$\begin{aligned} \tilde{P}_R(\theta, A) &= \int_A \left[\sum_{g' \in \mathcal{G}} \mathcal{P}_\vartheta^{(R)}(g'|g) \alpha_{ex}(\theta, \vartheta, g', g) \right] q_\theta(\vartheta|\theta) d\vartheta + \mathbb{I}(\theta \in A) \eta(\theta) \\ &\geq \int_A \left[\sum_{g' \in \mathcal{G}} \min \left\{ 1, \frac{\mathcal{P}_\vartheta^{(R)}(g'|g)}{\pi(g', \vartheta)} \right\} \pi(g', \vartheta) \alpha_{ex}(\theta, \vartheta, g', g) \right] q_\theta(\vartheta|\theta) d\vartheta \\ &\quad + \mathbb{I}(\theta \in A) \eta(\theta) > 0, \end{aligned}$$

where the last inequality comes from $\frac{\mathcal{P}_\vartheta^{(R)}(g'|g)}{\pi(g', \vartheta)} > 0$ for any $g' \in \mathcal{G}$ and $\vartheta \in \Theta$.

This proves that the statement is true when $s = 1$. By induction, we assume that it is true up to $s = n \geq 1$ and, for some $\theta \in \Theta$, choose $A \in \mathcal{B}(\Theta)$ such that $P^{(n+1)}(\theta, A) > 0$ and assume that

$$\int_\Theta \tilde{P}_R^{(n)}(\theta, d\vartheta) \tilde{P}_R(\vartheta, A) = 0.$$

This implies that $\tilde{P}_R(\vartheta, A) = 0$, $\tilde{P}_R^{(n)}(\theta, \cdot)$ -a.s.; by the induction assumption at $s = 1$, it follows that $P(\vartheta, A) = 0$, $\tilde{P}_R^{(n)}(\theta, \cdot)$ -a.s.

From this and the induction assumption at $s = n$, $P(\vartheta, A) = 0$, $P^{(n)}(\theta, \cdot)$ -a.s. (assume not, then $P(\vartheta, A) > 0$, $P^{(n)}(\theta, \cdot)$ -a.s. which by induction would imply $\tilde{P}_R(\vartheta, A) > 0$, which is a contradiction). The latter step contradicts $P^{(n+1)}(\theta, A) > 0$ and the result follows. *Q.E.D.*

The next step consists of proving that the transition kernel of the approximate algorithm $\tilde{P}_R(\theta, \vartheta)$ and the exact algorithm $P(\theta, \vartheta)$ are arbitrarily close for a large enough number of network simulations R . Formally, we prove a statement which is equivalent to proving convergence in total variation norm.⁴⁷

LEMMA 4—Convergence of the Exact and Approximate Transition Kernels: Let $\epsilon \in (0, 1]$. There exist a number of simulations $R_0 \in \mathbb{N}$ such that, for any function $\phi : \Theta \rightarrow [-1, 1]$ and any $R > R_0$,

$$|\tilde{P}_R \phi(\theta) - P \phi(\theta)| < 2\epsilon. \quad (35)$$

⁴⁷See Levin, Peres, and Wilmer (2008), Proposition 4.5, p. 49.

PROOF: The transition of the exchange algorithm is

$$P(\phi(\theta), \phi(\vartheta)) = \int_{\theta} \phi(\vartheta) \left[\sum_{g' \in \mathcal{G}} \pi(g', \vartheta) \alpha_{ex}(\theta, \vartheta, g', g) \right] q_{\theta}(\vartheta | \theta) d\vartheta \\ + \phi(\theta) \left[1 - \int_{\theta} \left[\sum_{g' \in \mathcal{G}} \pi(g', \vartheta) \alpha_{ex}(\theta, \vartheta, g', g) \right] q_{\theta}(\vartheta | \theta) d\vartheta \right],$$

while the transition kernel for the approximate algorithm is

$$\tilde{P}_R(\phi(\theta), \phi(\vartheta)) = \int_{\theta} \phi(\vartheta) \left[\sum_{g' \in \mathcal{G}} \mathcal{P}_{\vartheta}^{(R)}(g' | g) \alpha_{ex}(\theta, \vartheta, g', g) \right] q_{\theta}(\vartheta | \theta) d\vartheta \\ + \phi(\theta) \left[1 - \int_{\theta} \left[\sum_{g' \in \mathcal{G}} \mathcal{P}_{\vartheta}^{(R)}(g' | g) \alpha_{ex}(\theta, \vartheta, g', g) \right] q_{\theta}(\vartheta | \theta) d\vartheta \right],$$

and therefore the difference is

$$S = P(\phi(\theta), \phi(\vartheta)) - \tilde{P}_R(\phi(\theta), \phi(\vartheta)) \\ = \int_{\theta} \phi(\vartheta) \left[\sum_{g' \in \mathcal{G}} [\pi(g', \vartheta) - \mathcal{P}_{\vartheta}^{(R)}(g' | g)] \alpha_{ex}(\theta, \vartheta, g', g) \right] q_{\theta}(\vartheta | \theta) d\vartheta \\ - \phi(\theta) \int_{\theta} \left[\sum_{g' \in \mathcal{G}} [\pi(g', \vartheta) - \mathcal{P}_{\vartheta}^{(R)}(g' | g)] \alpha_{ex}(\theta, \vartheta, g', g) \right] q_{\theta}(\vartheta | \theta) d\vartheta.$$

Consider the quantity

$$S_0 = \int_{\theta} \left[\sum_{g' \in \mathcal{G}} [\pi(g', \vartheta) - \mathcal{P}_{\vartheta}^{(R)}(g' | g)] \alpha_{ex}(\theta, \vartheta, g', g) \right] q_{\theta}(\vartheta | \theta) d\vartheta \\ \leq \int_{\theta} \left[\sum_{g' \in \mathcal{G}} |\pi(g', \vartheta) - \mathcal{P}_{\vartheta}^{(R)}(g' | g)| \alpha_{ex}(\theta, \vartheta, g', g) \right] q_{\theta}(\vartheta | \theta) d\vartheta,$$

and since $\alpha_{ex}(\theta, \vartheta, g', g) \leq 1$ for any $(\theta, \vartheta) \in \Theta \times \Theta$ and $(g', g) \in \mathcal{G} \times \mathcal{G}$, we have

$$S_0 \leq \int_{\theta} \left[\sum_{g' \in \mathcal{G}} |\pi(g', \vartheta) - \mathcal{P}_{\vartheta}^{(R)}(g' | g)| \right] q_{\theta}(\vartheta | \theta) d\vartheta \\ = \int_{\theta} \left[2 \sup_{g' \in \mathcal{G}} |\pi(g', \vartheta) - \mathcal{P}_{\vartheta}^{(R)}(g' | g)| \right] q_{\theta}(\vartheta | \theta) d\vartheta.$$

The convergence of the network simulation algorithm implies that for any $\varepsilon > 0$, there exists an $R_0(\vartheta, \varepsilon) \in \mathbb{N}$ such that, for any $R > R_0(\vartheta, \varepsilon)$ and for any $g \in \mathcal{G}$,

$$2 \sup_{g' \in \mathcal{G}} |\pi(g', \vartheta) - \mathcal{P}_{\vartheta}^{(R)}(g' | g)| \leq \varepsilon.$$

Pick $R_0(\varepsilon) = \max_{\vartheta \in \Theta} \{R_0(\vartheta, \varepsilon)\}$. Then, for any $\varepsilon \in (0, 1]$, there is an $R_0(\varepsilon) \in \mathbb{N}$ such that, for any $R > R_0(\varepsilon)$ and for any $g \in \mathcal{G}$,

$$S_0 \leq \int_{\Theta} \varepsilon q_{\theta}(\vartheta | \theta) d\vartheta = \varepsilon.$$

This implies that $|S| \leq |2S_0| = 2\varepsilon$.

Q.E.D.

The next theorem is the main result for the convergence. It states that the approximate exchange algorithm converges to the correct posterior distribution, provided that the numbers of network simulations and parameter samples are big enough.

B.2.2. Proof of Theorem 6

PROOF: The main idea is to decompose the total variation in two components:

$$\begin{aligned} \|\tilde{P}_R^{(s)}(\theta_0, \cdot) - p(\cdot | g, X)\|_{\text{TV}} &= \|\tilde{P}_R^{(s)}(\theta_0, \cdot) - P^{(s)}(\theta_0, \cdot) + P^{(s)}(\theta_0, \cdot) - p(\cdot | g, X)\|_{\text{TV}} \\ &\leq \|\tilde{P}_R^{(s)}(\theta_0, \cdot) - P^{(s)}(\theta_0, \cdot)\|_{\text{TV}} + \|P^{(s)}(\theta_0, \cdot) - p(\cdot | g, X)\|_{\text{TV}}, \end{aligned}$$

and prove that each component converges. We will use the same idea, but rewrite the total variation in a more convenient form.⁴⁸ For any function $\phi : \Theta \rightarrow [-1, 1]$, we have

$$\begin{aligned} |\tilde{P}_R^{(s)} \phi(\theta_0) - p(\phi)| &= |\tilde{P}_R^{(s)} \phi(\theta_0) - P^{(s)} \phi(\theta_0) + P^{(s)} \phi(\theta_0) - p(\phi)| \\ &\leq |\tilde{P}_R^{(s)} \phi(\theta_0) - P^{(s)} \phi(\theta_0)| + |P^{(s)} \phi(\theta_0) - p(\phi)|. \end{aligned}$$

The second component converges because the exact exchange algorithm is ergodic, as stated in Lemma 2. For any $\varepsilon > 0$, there is a number of simulation steps $s(\theta_0, \varepsilon)$ such that, for any $s \geq s(\theta_0, \varepsilon)$,

$$|P^{(s)} \phi(\theta_0) - p(\phi)| \leq \varepsilon. \quad (36)$$

For the remaining of the proof, I will set $s_0 := s(\theta_0, \varepsilon)$. I use the telescoping sum decomposition in [Andrieu and Roberts \(2009\)](#) (p. 15, adapted from last formula)

$$\begin{aligned} |\tilde{P}_R^{(s_0)} \phi(\theta_0) - P^{(s_0)} \phi(\theta_0)| &= \left| \sum_{l=0}^{s_0-1} [P^{(l)} \tilde{P}_R^{(s_0-l)} \phi(\theta_0) - P^{(l+1)} \tilde{P}_R^{(s_0-(l+1))} \phi(\theta_0)] \right| \\ &= \left| \sum_{l=0}^{s_0-1} P^{(l)} (\tilde{P}_R - P) \tilde{P}_R^{(s_0-(l+1))} \phi(\theta_0) \right|. \end{aligned}$$

Now we can apply s_0 times the result of Lemma 4 (as in [Liang, Liu, and Carroll \(2010\)](#) and [Andrieu and Roberts \(2009\)](#)) to prove that there exists an $R_0(\theta_0, \varepsilon) \in \mathbb{N}$ such that, for any $R > R_0(\theta_0, \varepsilon)$,

$$|\tilde{P}_R^{(s_0)} \phi(\theta_0) - P^{(s_0)} \phi(\theta_0)| \leq 2s_0 \varepsilon, \quad (37)$$

this implies

$$|\tilde{P}_R^{(s)} \phi(\theta_0) - p(\phi)| \leq (2s_0 + 1) \varepsilon. \quad (38)$$

We conclude the proof by choosing $\varepsilon = \epsilon / (2s_0 + 1)$.

⁴⁸See [Levin, Peres, and Wilmer \(2008\)](#), proposition 4.5, p. 49.

This proves that the approximate exchange algorithm is ergodic, therefore the law of large number holds, and the second part of the theorem is proven. *Q.E.D.*

REFERENCES

- ANDRIEU, C., AND G. O. ROBERTS (2009): “The Pseudo-Marginal Approach for Efficient Monte Carlo Computations,” *Annals of Statistics*, 37 (2), 697–725. [8,12]
- DIACONIS, P., AND D. STROOCK (1991): “Geometric Bounds for Eigenvalues of Markov Chains,” *Annals of Applied Probability*, 1 (1), 36–61. [4]
- GELMAN, A., J. CARLIN, H. STERN, AND D. RUBIN (2003): *Bayesian Data Analysis, Second Edition*. Chapman & Hall/CRC. [8]
- GELMAN, A., G. O. ROBERTS, AND W. R. GILKS (1996): “Efficient Metropolis Jumping Rules,” *Bayesian Statistics*, 5, 599–608. [3]
- LEVIN, D. A., Y. PERES, AND E. L. WILMER (2008): *Markov Chains and Mixing Times*. American Mathematical Society. [4,10,12]
- LIANG, F. (2010): “A Double Metropolis-Hastings Sampler for Spatial Models With Intractable Normalizing Constants,” *Journal of Statistical Computing and Simulation*, 80 (9), 1007–1022. [4,8]
- LIANG, F., C. LIU, AND R. J. CARROLL (2010): *Advanced Markov Chain Monte Carlo Methods: Learning From Past Samples*. Wiley. [8,12]
- MONDERER, D., AND L. SHAPLEY (1996): “Potential Games,” *Games and Economic Behavior*, 14, 124–143. [1, 2]
- MURRAY, I. A., Z. GHAHRAMANI, AND D. J. C. MACKAY (2006): “MCMC for Doubly-Intractable Distributions,” *Uncertainty in Artificial Intelligence*. [7]
- ROBERT, C. P., AND G. CASELLA (2005): *Monte Carlo Statistical Methods (Springer Texts in Statistics)*. New York: Springer. [8]

Carey Business School, Johns Hopkins University, 100 International Drive, Baltimore, MD 21202, U.S.A.; angelo.mele@jhu.edu.

Co-editor Matthew O. Jackson handled this manuscript.

Manuscript received 4 November, 2011; final version accepted 31 October, 2016; available online 18 January, 2017.