

1 STATIONARY SOCIAL LEARNING IN A CHANGING ENVIRONMENT 1

2
3 RAPHAËL LEVY 3

4 Department of Economics and Decision Sciences, HEC Paris 4

5
6 MARCIN PEŃSKI 6

7 Department of Economics, University of Toronto 7

8
9 NICOLAS VIEILLE 8

10 Department of Economics and Decision Sciences, HEC Paris 9

11 We consider social learning in a changing world. With changing states, soci- 11
12 eties can be responsive only if agents regularly act upon fresh information, which 12
13 significantly limits the value of observational learning. When the state is close 13
14 to persistent, a consensus whereby most agents choose the same action typically 14
15 emerges. However, the consensus action is not perfectly correlated with the state, 15
16 because societies exhibit inertia following state changes. When signals are precise 16
17 enough, learning is incomplete, even if agents draw large samples of past actions, 17
18 as actions then become too correlated within samples, thereby reducing informa- 18
19 tiveness and welfare. 19

20 1. INTRODUCTION 20

21 The literature on social learning has extensively studied the extent to which agents learn 21
22 from others' actions. In particular, it has been quite successful at understanding the possi- 22
23 ble emergence of informational cascades, and conditions under which the consensus that 23
24 eventually forms over time is correct (see [Bikhchandani et al. \(2021\)](#) for a recent survey). 24
25 However, little attention has been drawn to the possibility that the underlying state of na- 25
26 ture is changing. 26

27
28

Raphaël Levy: levyr@hec.fr 28

29 Marcin Peński: marcin.peski@utoronto.ca 29

30 Nicolas Vieille: vieille@hec.fr 30

ture might change over time.¹ Still, in several applications, e.g., technology adoption or investment decisions, the optimal course of action is likely to evolve, raising the question of whether social learning then efficiently aggregates information.

The possibility of state changes provides new insights both from applied and theoretical perspectives. For instance, the dynamics of learning may shed light on how societies react to changes in the environment, and on how a dominant consensus may be replaced by a new one.² From a theoretical perspective, the possibility of state changes creates a tension between information aggregation and responsiveness to change. Indeed, while efficient aggregation supposes that some agents safely rely on their peers to act, it also requires that society reacts swiftly to a change in the environment, ruling out informational cascades. The goal of this paper is to evaluate how this tension shapes equilibrium welfare.

To address this question, we analyze steady-state equilibria in a model where (a) the state of nature follows a Markov chain and (b) in each period, a continuum of short-lived agents draw a finite sample of past actions and have access to a (possibly costly) informative signal. Though unnecessary for our results, allowing for costly information makes social learning even more desirable. Indeed, the potential welfare gains from learning from others result not only from better-informed decisions, but also from savings on information costs.

Two forces drive equilibrium welfare at a steady state. The first one is reminiscent of the Grossman-Stiglitz paradox: actions cannot be too informative about the state for otherwise agents would have no incentive to acquire information. With changing states, this logic imposes that some fresh information flows in every period at the steady state, and thus puts a limit on equilibrium welfare. When agents sample at least two actions, a second (countervailing) force comes into play. In that case, some agents will sample conflicting evidence, and rely entirely on their signals, while others will obtain unambiguous evidence

¹Exceptions are [Moscarini, Ottaviani and Smith \(1998\)](#), [Acemoglu, Nedic and Ozdaglar \(2008\)](#), [Frongillo, Schoenebeck and Tamuz \(2011\)](#), [Huang \(2022\)](#), [Dasaratha, Golub and Hak \(2023\)](#). See Section 6 for more detail.

²Examples of a change in the dominant technology abound, ranging from the “war of the currents” in the late 19th century, the “quartz crisis” in watchmaking in the 1970s to Facebook overtaking MySpace as the dominant social network.

1 allowing them to possibly free-ride on the information acquired by the former. This creates 1
2 a inter-temporal externality across samples which can be expected to bring welfare gains. 2
3 This is exactly what happens in [Banerjee and Fudenberg \(2004\)](#), who show in a fixed-state 3
4 model that learning is eventually complete when agents sample as few as two actions. In 4
5 sharp contrast, our chief finding is that allowing for even arbitrarily rare changes in the state 5
6 typically results in incomplete learning. 6

7 We first analyze the case where agents sample at most two actions. In any equilibrium, 7
8 *all* agents must acquire (or make use of) information with positive probability, regardless 8
9 of their samples. Otherwise, indeed, the forces of imitation are so strong that all agents 9
10 within and across periods eventually play the same action. In a changing world, such an 10
11 uninformative, irreversible consensus is ruled out in equilibrium. Therefore, learning is 11
12 incomplete: agents never entirely rely on their peers, even when the state is arbitrarily 12
13 persistent. Not only observing two actions no longer ensures complete learning, but welfare 13
14 is typically worse than when only one action is sampled. 14
15

16 For larger samples, we focus on the case where the state is highly persistent. First, we 16
17 show that a consensus prevails. That is, in a steady-state equilibrium, most likely most 17
18 agents play the same action. However, this consensus must be fluctuating, and the popula- 18
19 tion oscillates over time following state changes. The dynamics following a state change is 19
20 characterized by two phases. Agents first stick to the current consensus unless their sample 20
21 conveys somewhat mixed evidence. But such conflicting evidence is unlikely in a society 21
22 where one action dominates, and there may be significant inertia in moving away from an 22
23 established consensus. Once the population displays some minimal dissent, the fraction of 23
24 agents acting against the old consensus quickly takes off, and the population snowballs to- 24
25 wards a new consensus. The efficiency of learning reflects responsiveness, that is, how long 25
26 it takes for society to escape an obsolete consensus when the state changes. 26

27 We next show that, when signals are binary, precise enough, and not too costly, there 27
28 exists an equilibrium in which welfare is the same as in no-social-learning benchmark 28
29 where agents observe no past action. When signals get more precise, agents are more likely 29
30 to play the right action when acquiring information, and both the convergence towards a 30

correct consensus and away from a wrong consensus get faster. However, it turns out that the *relative* speed of convergence towards a correct consensus increases. As for samples of size two, agents observing even unanimous samples must then acquire information to make sure that society remains responsive to change and does not get stuck in an irreversible consensus. Intuitively, when signals are precise, the actions of agents acquiring information are highly correlated to the state, hence among themselves. Actions are less diverse, which curtails information aggregation and, thereby, welfare.

To prove this result, we exploit time-invariance equations for measures to provide bounds on the beliefs for non-unanimous sample compositions. Intuitively, when state changes are rare, equilibrium beliefs can be approximated in terms of time-average values of the dynamical systems that describe the evolution of the population’s behavior. Perhaps surprisingly, these belief estimates point to the presence of belief reversal: seeing one dissenting action within an otherwise unanimous sample should be taken as evidence that the minority action is the correct one.³

The paper is organized as follows. We introduce the model in Section 2. Section 3 is devoted to small samples, and Section 4 provides results for larger samples in the persistent limit case. Section 5 addresses robustness issues and extensions. We discuss the relation to the literature in Section 6, and conclude in Section 7.

2. THE MODEL

2.1. States, actions and payoffs

We consider a social learning model in discrete time with an evolving, binary state of nature $\theta \in \Theta := \{0, 1\}$. In each period, there is a continuum of short-lived agents who choose an action from the action set $A := \{0, 1\}$ and obtain a utility of one when their action matches the current state, and of zero otherwise.

³In the online appendix, we further leverage this approach to numerically investigate equilibrium behavior and welfare in the case of samples of size three. For signals of low precision, equilibrium welfare may be higher than in the no-social-learning case, unlike for precise signals. Yet, learning is always incomplete, in line with the main message of the paper.

Successive states (θ_t) follow a symmetric Markov chain over Θ . The parameter $\lambda := \mathbf{P}(\theta_{t+1} \neq \theta \mid \theta_t = \theta)$ captures the degree of persistency. States are i.i.d. if $\lambda = \frac{1}{2}$, and fully persistent if $\lambda = 0$. We assume that $\lambda \in (0, \frac{1}{2})$: the state is persistent, but not fully.

2.2. Timing, sampling and signals

At each date t , events unfold as follows. Each new-born agent (i) first observes a random sample of n past actions, (ii) next decides whether or not to acquire additional information about the current state θ_t at cost $c \geq 0$, (iii) finally picks an action $a \in A$.

We assume that sampled actions are drawn from the pool of actions played in the previous period, in proportion to their prevalence in the population (proportional sampling). That is, the sample composition at date t , measured by the count of ones, follows a binomial distribution $B(n, x_{t-1})$, where x_{t-1} is the fraction of agents playing action 1 in period $t - 1$. Samples are independent across agents and private.

The additional information available to agents consists of a private signal that is independent across agents conditional on the current state. We let q denote the posterior belief assigned to $\theta = 1$ given the signal under a uniform prior, and refer to q as a *private belief*. We denote by H_θ the right-continuous cdf of q in state θ . Signal distributions are assumed symmetric across states.⁴ That is, the distribution of the posterior probability assigned to θ conditional on the state being θ is the same for both states. This corresponds to $H_0(q)_- = 1 - H_1(1 - q)$ for each $q \in [0, 1]$. We rule out uninformative signals, and assume throughout that $H_1(q) < H_0(q)$ for some $q \in (0, 1)$. Finally, we denote by \bar{q} the supremum of the support of the unconditional distribution $H := \frac{1}{2}H_0 + \frac{1}{2}H_1$. By symmetry, the infimum is $1 - \bar{q}$. Following the usual terminology, signals are *unbounded* if $\bar{q} = 1$, and *bounded* if $\bar{q} < 1$.

⁴The asymmetric case is discussed in Section 5.2.

2.3. Equilibrium concept

We focus on equilibrium steady states in which all agents across and within periods use the same decision rule σ . The equilibrium notion requires that σ is optimal given beliefs, and that beliefs are derived from the invariant joint distribution of states and samples induced by σ . In addition, we restrict attention to symmetric equilibria, i.e., we require that the equilibrium is unchanged when relabelling actions and states so that the decisions given a sample $n - k$ are the mirror images of those made with sample k . A formal definition of equilibrium steady states is given in Section 4.1, together with an existence result. At this stage, we simply denote by p_k the (interim) probability that the current state is $\theta = 1$ conditional on seeing a sample composed of $k \in \{0, \dots, n\}$ ones.

2.4. Information acquisition

Consider an agent who holds an interim belief p and contemplates acquiring information. Upon acquiring information and receiving a signal inducing a private belief q , the agent chooses action 1 whenever the more likely state is $\theta = 1$, i.e., if $pq \geq (1 - p)(1 - q) \Leftrightarrow q \geq 1 - p$, with indifference if $q = 1 - p$.

Accordingly, the probability of playing the correct action $a = \theta$ is given by

$$v(p) := p(1 - H_1(1 - p)) + (1 - p)H_0(1 - p). \quad (1)$$

The function v is convex, increasing on $[\frac{1}{2}, 1]$ and symmetric: $v(p) = v(1 - p)$ for all p .

Instead, when information is not acquired, the agent's action matches the state with probability $u(p) := \max(p, 1 - p)$.

Since more information cannot hurt, $u(p) \leq v(p)$ for all p ; besides, the net value of acquiring information $v(p) - u(p)$ is maximal when $p = \frac{1}{2}$. If $c > v(\frac{1}{2}) - u(\frac{1}{2})$, agents never acquire information and samples are not informative. We rule out this case and assume throughout:


ASSUMPTION 1: $v(\frac{1}{2}) - u(\frac{1}{2}) \geq c$.

1 These properties imply the existence of a unique $\hat{p} \in [\frac{1}{2}, 1]$ such that $v(p) - c > u(p)$ iff
 2 $p \in (1 - \hat{p}, \hat{p})$. As our analysis highlights, what ultimately matters is whether $\hat{p} < 1$ or not.

3 When signals are bounded ($\bar{q} < 1$), one has $v(p) = u(p)$ for $p > \bar{q}$, hence $\hat{p} < 1$ irrespec-
 4 tive of whether $c = 0$ or $c > 0$. When signals are unbounded, $v(p) > u(p)$ for all $p \in (0, 1)$,
 5 hence $\hat{p} = 1$ if $c = 0$ and $\hat{p} < 1$ if $c > 0$. In what follows, we maintain the assumption that
 6 $\hat{p} < 1$ (the richer case), and defer the discussion of the easier case $\hat{p} = 1$ to Section 5.1.

7 ASSUMPTION 2: $\hat{p} < 1$.

8
 9 Figures 1 and 2 illustrate two typical cases. In Figure 1, signals are unbounded and
 10 $c > 0$; in Figure 2, signals are binary with precision $\frac{1}{2} < \pi < 1$ (bounded). In that case,
 11 private beliefs are either $1 - \pi$ or π , and $v(p) = \max(p, 1 - p, \pi)$, implying $\hat{p} = \pi - c$.



12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
FIGURE 1.—Signals with unbounded strength




FIGURE 2.—Binary signals with precision π

We conclude this section with a property that any equilibrium must satisfy, namely, there is no steady-state cascade.

LEMMA 1: *In any equilibrium steady state, one has $1 - \hat{p} \leq p_k \leq \hat{p}$ for some k .*

PROOF: Assume instead that for some equilibrium and for each k , one has either $p_k < 1 - \hat{p}$ or $p_k > \hat{p}$. Signals are never used, hence samples are uninformative at the steady state and therefore, $p_k = \frac{1}{2}$ for each k . A contradiction. *Q.E.D.*

Lemma 1 imposes that private signals must be used with positive probability following at least one sample realization in any steady-state equilibrium. Still, one may have $p_k \notin$

[1 - \hat{p} , \hat{p}] for *some* sample compositions. Upon observing such samples, agents play their perceived best action without using further information.

3. SMALL SAMPLES

We discuss here small sample sizes ($n \leq 2$). The case $n = 0$ where agents do not sample serves as a no-social-learning benchmark. In that case, the belief of all agents at a steady state is given by the invariant distribution of (θ_t) , which is uniform over Θ . Agents acquire information, and obtain an expected payoff of $v(\frac{1}{2}) - c$.

3.1. Samples of size $n = 1$

We assume here that $n = 1$. Equilibrium symmetry implies that $p_1 = 1 - p_0$ and that all agents acquire information with the same probability β , with $\beta > 0$ by Lemma 1. We set $\lambda_* := \frac{c}{2(\hat{p}+c)-1} < \frac{1}{2}$. Note that $\lambda_* = 0$ when $c = 0$.

PROPOSITION 1: *If $n = 1$, there is a unique equilibrium steady state:*

- If $\lambda < \lambda_*$, one has $p_1 = \hat{p}$ and $\beta = \lambda \frac{2\hat{p} - 1}{c(1 - 2\lambda)} \in (0, 1)$.
- If $\lambda \geq \lambda_*$, one has $p_1 \in (1 - \hat{p}, \hat{p})$ and $\beta = 1$.

PROOF: Consider an agent A in period t who samples the action a_{t-1} of some player B. In the steady state, both A and B hold either an interim belief p_1 or $p_0 = 1 - p_1$, where p_1 obeys the following equation:

$$\begin{aligned} p_1 &= \mathbf{P}(\theta_t = 1 \mid a_{t-1} = 1) \\ &= (1 - \lambda)\mathbf{P}(\theta_{t-1} = 1 \mid a_{t-1} = 1) + \lambda\mathbf{P}(\theta_{t-1} = 0 \mid a_{t-1} = 1). \end{aligned} \quad (2)$$

Either, with probability β , B acquired information and then played the right action with probability $v(p_1)$, or did not, and matched the state with probability $u(p_1)$.

If $0 < \beta < 1$, B's indifference condition imposes $p_1 = \hat{p}$. One therefore derives $\mathbf{P}(\theta_{t-1} = 1 \mid a_{t-1} = 1) = \beta v(\hat{p}) + (1 - \beta)u(\hat{p}) = \hat{p} + \beta c$. Substituting into (2) yields $\hat{p} = \lambda + (1 - 2\lambda)(\hat{p} + \beta c)$, hence $\beta = \lambda \frac{2\hat{p} - 1}{c(1 - 2\lambda)}$. $\beta < 1$ then requires $\lambda < \lambda_*$.

If $\beta = 1$, one has $\mathbf{P}(\theta_{t-1} = 1 \mid a_{t-1} = 1) = v(p_1)$ and (2) now reads

$$p_1 = (1 - \lambda)v(p_1) + \lambda(1 - v(p_1)). \quad (3)$$

Since $v'(p_1) < 1$ for $p_1 \leq \hat{p}$ and since $v(\frac{1}{2}) - c > \frac{1}{2}$, (3) has a (unique) solution in $[\frac{1}{2}, \hat{p}]$ if and only if $\lambda \geq \lambda_*$. *Q.E.D.*

When the state changes frequently, *past* actions cannot possibly be very informative about the *current* state, and information is acquired with probability 1. As the state gets more persistent, past actions potentially become informative, and β decreases. In the persistent limit $\lambda \rightarrow 0$, agents acquire information with vanishing probability, hence most likely replicate the action they sample. The equilibrium payoff increases from $v(\frac{1}{2}) - c$ to \hat{p} as λ decreases from the i.i.d. case to the persistent limit. More generally, the equilibrium payoff, hence welfare, is given as follows.

COROLLARY 1: *The equilibrium welfare is $v(p_1) - c$ for $\lambda \geq \lambda_*$, where p_1 is the solution of (3), and $p_1 = \hat{p}$ for $\lambda < \lambda_*$.*

For binary signals with precision π , the equilibrium welfare is $\pi - c = \hat{p}$, for each $\lambda > 0$.

3.2. Samples of size $n = 2$

When $n = 1$, private signals are always interim valuable in equilibrium, which limits the informativeness of past actions. The situation is quite different with larger samples. With $n > 1$, the efficiency of social learning can be improved if agents with some sample k generate enough information that (future) agents with a different sample k' find it optimal to herd – that is, if there exists (k, k') such that $1 - \hat{p} \leq p_k \leq \hat{p} < p_{k'}$. Ultimately, the equilibrium welfare is determined by the magnitude of such information externalities across samples.

Let us look at how this discussion applies to the case $n = 2$. In this case, $p_1 = \frac{1}{2}$, by symmetry: agents who sample conflicting actions are confused and acquire information. The key question is whether the information produced by these agents is enough to ensure that other agents can herd, i.e., $p_2 > \hat{p}$. Casual intuition suggests that this should be the case when the state is sufficiently persistent. But, remarkably, this intuition is incorrect.

PROPOSITION 2: *In any equilibrium, one has $p_2 \in [1 - \hat{p}, \hat{p}]$.*

Proposition 2 implies that agents are *always* willing to acquire information when $c > 0$. For $c = 0$, it implies that it can *never* be strictly optimal to ignore one's signal.

The logic works as follows. Assume that $p_2 > \hat{p}$, so that agents acquire information when sampling conflicting evidence only. By acquiring information, these agents are instrumental in moving towards a correct consensus. Sooner or later, society will reach such a consensus, at which point there will be too few agents sampling mixed evidence, and the population will no longer be responsive to changes in the state.

The complete proof of Proposition 2 is in the Appendix. We provide a sketch below.

PROOF SKETCH: We argue by contradiction and assume that $p_2 > \hat{p}$ in some equilibrium. Since $p_1 = \frac{1}{2}$, agents with a balanced sample acquire information, and choose action 1 if their private belief exceeds $\frac{1}{2}$, which has probability $\phi_\theta := 1 - H_\theta(\frac{1}{2})$ in state θ .

Denoting by x_{t-1} the fraction of agents playing action 1 in period $t - 1$, the probability that a generic agent in period t plays action 1 is thus given by $x_t = \bar{g}_\theta(x_{t-1})$, where

$$\bar{g}_\theta(x) := x^2 + 2x(1-x)\phi_\theta.$$

Since $\phi_1 > \frac{1}{2} > \phi_0$, one has $\bar{g}_1(x) > x > \bar{g}_0(x)$ for each $x \in (0, 1)$: the popularity x_t of action 1 increases over time when $\theta_t = 1$, and decreases otherwise, as shown in Figure 3.

For x close to 0, the ratio $\bar{g}_\theta(x)/x$ is approximately equal to $2\phi_\theta$. Thus, as long as x_t is close to zero, $\ln x_t$ increases by $\ln 2\phi_1$ when $\theta_t = 1$, and decreases by $\ln 2\phi_0$ when $\theta_t = 0$. Since $4\phi_0\phi_1 = 4\phi_1(1-\phi_1) < 1$, one has $\ln 2\phi_1 < -\ln 2\phi_0$: step sizes are higher when $\ln x_t$ decreases. This means that (x_t) converges faster *towards* a consensus on action 0 when $\theta_t = 0$ than it moves *away* from this consensus when $\theta_t = 1$. Consequently, whenever (x_t) approaches either 0 or 1, there is a positive probability that the population will never bounce away from this consensus, even following state changes.

Combined with the observation, obvious from Figure 3, that (x_t) cannot stay bounded away from 0 and 1 indefinitely, this implies that the sequence (x_t) converges to either 0 or 1, almost surely: the population converges to a *permanent* consensus, although the state of




FIGURE 3.—Population dynamics

nature keeps changing. Thus, past actions are fully uninformative at the steady state hence $p_2 = \frac{1}{2}$. A contradiction. *Q.E.D.*

COROLLARY 2: *When signals are binary, the equilibrium welfare is \hat{p} for all $n = 0, 1, 2$, all $\lambda > 0$ and all $c \geq 0$. When H_θ has support $[1 - \bar{q}, \bar{q}]$, the equilibrium welfare is strictly lower for $n = 2$ than for $n = 1$ if $\lambda \leq \lambda_*$.*

PROOF: By Proposition 2, acquiring information is always valuable with $n = 2$, so an agent obtains a payoff $v(p_k) - c$ for any sample realization k , where $p_k \in [1 - \hat{p}, \hat{p}]$. This implies that welfare is no larger than \hat{p} , that is, the welfare when $n = 1$ in the case $\lambda \leq \lambda_*$.

With binary signals with precision π , $v(p) = \pi$ for all $p \in [1 - \hat{p}, \hat{p}]$, and the equilibrium payoff is then $\pi - c = \hat{p}$ for every $\lambda > 0$ and $c \geq 0$, like in the cases $n = 0$ and $n = 1$.

When H_θ has support $[1 - \bar{q}, \bar{q}]$, v is strictly convex on $[1 - \hat{p}, \hat{p}]$, and $v(p_1) - c = v(\frac{1}{2}) - c < \hat{p}$. *Q.E.D.*

When signals are binary, social learning arises in equilibrium (some agents do herd), yet there is no welfare gain over the benchmark case $n = 0$ where information is always acquired. With richer signals, the equilibrium welfare is non-monotonic for small λ : it increases from $n = 0$ to $n = 1$ and decreases from $n = 1$ to $n = 2$. This is all the more surpris-

ing as this holds when the state is highly persistent, that is, when the logic of observational learning acts most forcefully.

3.3. Comparison to fixed state models

We here relate these findings to the literature with a fixed state and known calendar time. The relevant comparison is [Banerjee and Fudenberg \(2004\)](#) (henceforth BF), who assume that in each period, a continuum of short-lived agents draw a random sample of past actions and observe a signal for free.⁵ If $n = 1$, the analysis of BF implies the existence of a continuum of steady states. In any such steady state, the fraction of agents who play the correct action is constant over time, and equal to $p_1 \geq \hat{p}$. In each period, all agents ignore their signals, and replicate the action they sample.⁶ This multiplicity is ruled out with changing states, since it cannot be that agents systematically ignore their signals.

If $n = 2$, our results sharply contrast with BF. For $n \geq 2$, BF shows that, under minimal assumptions on signals (that are satisfied in our setup), learning is eventually complete: actions converge to the correct one. As soon as the state may change, our analysis instead shows that equilibrium payoffs do not exceed \hat{p} , even as $\lambda \rightarrow 0$. Accordingly, allowing for a changing state has a strong negative impact. In addition, the comparison between samples of size 1 and 2 suggests that, while observing more actions compensates for a limited signal quality when the state is fixed, this may exacerbate the inefficiency in a changing world.

4. THE GENERAL CASE: EQUILIBRIUM ANALYSIS

The discussion on small samples illustrates a key complication. When only one action is sampled, the interim belief at date t involves only the *expected* value of x_{t-1} in each state, reflecting how often on average past agents play the correct action. This allows for a closed form analysis. As soon as two actions are sampled, the interim belief p_k also involves the

⁵Although BF considers free signals while we allow for costly signals, this distinction is irrelevant: on the one hand, our results apply to free signals without loss; on the other hand, the results of BF still hold with costly signals (we provide a proof in the supplementary material).

⁶Notice that since there is a continuum of agents, not all agents play the same action.

1 correlation of actions within samples.⁷ Formally, the belief likelihood ratio is

$$\frac{\mathbf{P}(\theta_{t-1} = 1 \mid k)}{\mathbf{P}(\theta_{t-1} = 0 \mid k)} = \frac{\int_0^1 x^k (1-x)^{n-k} d\mu_1(x)}{\int_0^1 x^k (1-x)^{n-k} d\mu_0(x)} \quad (4)$$

2
3
4
5
6 where μ_θ is the distribution of x_{t-1} in state θ , so that the formula for p_k involves all l -th
7 moments of x_{t-1} for $l \leq n$; in addition, the evolution of x_t^l over time involves even higher
8 powers of x_t , as can be checked.

9 Consequently, the equilibrium conditions involve the entire joint steady-state distribution
10 of (θ_t, x_t) . This distribution is a complex object, leaving little hope for a tractable analysis.⁸
11 Accordingly, we focus on the persistent limit $\lambda \rightarrow 0$, which we view as the most relevant
12 case for a comparison with the usual fixed-state setup.

13 In Section 4.1, we provide a formal definition of strategies and equilibrium steady-states.
14 In Section 4.2, we prove that the population is (asymptotically) always in consensus. Sec-
15 tion 4.3 discusses welfare and the extent to which the prevailing consensus is correct.

16 17 4.1. Strategies and equilibrium

18 A strategy specifies whether or not to acquire information (if $c > 0$) and which action
19 to choose. These decisions depend on the composition of one's sample, and (whenever
20 relevant) on the signal. A strategy is thus a pair $\sigma = (\beta, \alpha)$ of (measurable) maps, with
21 $\beta : \{0, \dots, n\} \rightarrow [0, 1]$ and $\alpha : \{0, \dots, n\} \times [0, 1] \rightarrow \Delta(\{0, 1\})$, with the understanding that
22 $\beta(k)$ is the probability of acquiring information upon observing a sample composed of k
23 ones, and $\alpha(k, q)$ is the probability of playing action 1 upon drawing sample k and ob-
24 serving a private belief $q \in [0, 1]$. Not acquiring information is informationally equivalent
25 to drawing an uninformative signal $q = \frac{1}{2}$ for sure: an agent with sample k who does not
26 acquire information thus plays 1 with probability $\alpha(k, \frac{1}{2})$.

27
28 ⁷Actions sampled in period t are independent conditional on x_{t-1} , but correlated *ex ante*.

29 ⁸Even in the much simpler case where states are i.i.d. and $n = 1$, extensive work has focused on the properties
30 of the steady-state distribution of x . See, e.g. Solomyak (1995) or Bhattacharya and Majumdar (2007).

Conditional on the state being θ , an agent sampling k thus plays action 1 with probability

$$\phi_\theta(k) := \beta(k) \int_0^1 \alpha(k, q) dH_\theta(q) + (1 - \beta(k)) \alpha(k, \frac{1}{2}). \quad (5)$$

The sample composition at date t follows a Binomial distribution $B(n, x_{t-1})$. It follows that the fraction of agents choosing action 1 in period t is

$$x_t = g_{\theta_t}(x_{t-1}) := \sum_{k=0}^n \binom{n}{k} x_{t-1}^k (1 - x_{t-1})^{n-k} \phi_{\theta_t}(k) \quad (6)$$

For fixed σ , the pair (θ_t, x_t) follows a Markov chain over $\Theta \times [0, 1]$. Over time, θ_t evolves independently of x_t , and $x_t = g_{\theta_t}(x_{t-1})$.

An *equilibrium steady state* is a pair (μ, σ) where $\mu \in \Delta(\Theta \times [0, 1])$ is an invariant measure for (θ_t, x_t) , and σ is optimal given μ . The optimality condition on σ reads as **C1** and **C2** below:

C1 $\beta(k) = 1$ if $p_k \in (1 - \hat{p}, \hat{p})$ and $\beta(k) = 0$ if $p_k \notin [1 - \hat{p}, \hat{p}]$;⁹

C2 $\alpha(k, q) = 1$ if $q > 1 - p_k$ and $\alpha(k, q) = 0$ if $q < 1 - p_k$.

We recall that the interim belief $p_k = \mathbf{P}(\theta_t = 1 \mid k)$ is the belief on the *current* state. It is related to the belief on the *previous* state through the equality $p_k = (1 - \lambda)\mathbf{P}(\theta_{t-1} = 1 \mid k) + \lambda\mathbf{P}(\theta_{t-1} = 0 \mid k)$. The belief $\mathbf{P}(\theta_{t-1} = 1 \mid k)$ is itself related to the steady-state distribution μ through (4).

The condition that μ is an invariant distribution for (θ_t, x_t) reads

C3 $\mu(\theta, X) = (1 - \lambda)\mu(\theta, g_\theta^{-1}(X)) + \lambda\mu(1 - \theta, g_\theta^{-1}(X))$ for all measurable $X \subset [0, 1]$.

Since we focus on symmetric equilibria, we require in addition that μ and σ treat the two states and actions symmetrically. Formally:

C4 $\beta(k) = \beta(n - k)$ and $\alpha(k, q) = 1 - \alpha(n - k, 1 - q)$ for each k and q .

C5 μ is invariant under the transformation $(\theta, x) \mapsto (1 - \theta, 1 - x)$.

⁹If $c = 0$, condition **C1** can be omitted. Indeed, if $p_k > \hat{p}$, it is optimal to play $a = 1$ for all q so that $\alpha(k, q) = 1$. For such k , $\phi_\theta(k) = 1$ irrespective of $\beta(k)$. Similarly, when $p_k < 1 - \hat{p}$, $\phi_\theta(k) = 0$ for any $\beta(k)$.

We denote by $G(\lambda)$ the game with transition parameter λ . The proof of Theorem 1 below is in Appendix B.

THEOREM 1: *The game $G(\lambda)$ has a symmetric equilibrium steady state.*

4.2. Aggregate behavior: a consensus result

We first derive a general result on the aggregate behavior in the population, and prove that the actions of agents become highly correlated as the state gets close to persistent.

THEOREM 2: *Let $n \geq 2$ and let $(\mu_\lambda, \sigma_\lambda)$ be any equilibrium steady state of $G(\lambda)$, for $\lambda > 0$. As $\lambda \rightarrow 0$, the marginal of μ_λ over $x \in [0, 1]$ (weakly) converges to the uniform distribution over the two-point set $\{0, 1\}$.*

According to Theorem 2, most likely most agents play the same action in any given period. This consensus result implies in turn that most likely most agents draw a unanimous sample $k \in \{0, n\}$ consisting only of zeroes or of ones.

PROOF: Let λ and an equilibrium (μ, σ) of $G(\lambda)$ be given, and denote by κ^* the (steady-state) average fraction of agents whose action matches the state. κ^* is weakly higher than the equilibrium payoff w^* because information acquisition costs are not accounted for.

Let us list the actions within a generic agent's sample in some random order $a^{(1)}, \dots, a^{(n)}$. One available strategy σ_1 is to simply imitate $a^{(1)}$. The strategy σ_1 would yield a payoff of κ^* if the state were fixed. Accounting for state transitions, and assuming $a^{(1)} = 1$ for concreteness, σ_1 yields

$$w(\sigma_1) := \mathbf{P}(\theta_t = 1 \mid a^{(1)} = 1) = (1 - \lambda)\kappa^* + \lambda(1 - \kappa^*) \geq \kappa^* - \lambda,$$

and thus, $w(\sigma_1) \geq w^* - \lambda$. Since no strategy improves upon w^* , this implies that the marginal gain of observing $a^{(2)}$ is at most λ .

In turn, this implies that $a^{(1)}$ and $a^{(2)}$ coincide with high probability when λ is small. Indeed, consider the alternative strategy σ_2 consisting in playing $a^{(1)}$ if the second action confirms the first one ($a^{(1)} = a^{(2)}$) and acquiring information otherwise ($a^{(1)} \neq a^{(2)}$). In

1 the latter case, the agent's belief is $1/2$, hence the agent's payoff conditional on $a^{(1)} \neq a^{(2)}$ 1
 2 is $v(1/2) - c$. Therefore, the payoff $w(\sigma_2)$ is a convex combination of $v(1/2) - c$ and of 2
 3 $\mathbf{P}(\theta_t = 1 \mid a^{(1)} = a^{(2)} = 1)$, where the weights are the conditional probabilities of $a^{(2)} = 0$ 3
 4 and of $a^{(2)} = 1$ given $a^{(1)} = 1$. 4

5 On the other hand, the martingale property of beliefs ensures that $\mathbf{P}(\theta_t = 1 \mid a^{(1)} = 1)$ 5
 6 (which is also $w(\sigma_1)$) is a convex combination of the beliefs $1/2$ and of $\mathbf{P}(\theta_t = 1 \mid a^{(1)} =$ 6
 7 $a^{(2)} = 1)$, with the *same* weights. Since $v(1/2) - c > u(1/2) = 1/2$, and since $w(\sigma_2) \leq$ 7
 8 $w^* \leq w(\sigma_1) + \lambda$, it follows that the probability $\mathbf{P}(a^{(2)} = 0 \mid a^{(1)} = 1)$ that $a^{(2)}$ contradicts 8
 9 $a^{(1)}$ is at most of the order of λ . 9

10 To conclude, recall that $a^{(1)}$ and $a^{(2)}$ are independent draws from a Bernoulli distribution 10
 11 with parameter x , where x is first drawn according to μ . Since $a^{(1)}$ and $a^{(2)}$ coincide with 11
 12 high probability, it must be that x is quite close to 0 or to 1, with high μ -probability. *Q.E.D.* 12

13 According to Theorem 2, the population is in consensus in a typical period, with x_t being 13
 14 close to either 0 or 1. At the same time, the consensus must evolve over time in response to 14
 15 changes in the state. This implies that the population alternates between the two consensus, 15
 16 and that the transition periods are vanishingly short relative to the time spent in consensus. 16
 17

18 Let us briefly elaborate on the transition dynamics, assuming $n = 2$ and $c > 0$ for con- 18
 19 creteness. We denote by $\beta := \beta(0) = \beta(2) > 0$ the equilibrium probability of acquiring 19
 20 information when sampling $k \in \{0, 2\}$. Theorem 2 implies that, as $\lambda \rightarrow 0$, fewer and fewer 20
 21 agents acquire information, hence $\beta \rightarrow 0$. For λ small, interim beliefs are $p_2 = 1 - p_0 = \hat{p}$ 21
 22 and $p_1 = \frac{1}{2}$. Since agents either acquire information or herd, it can be checked that x_t 22
 23 evolves according to $x_t = g_{\theta_t}(x_{t-1})$, with 23

$$24 \quad g_{\theta}(x) = x^2 + 2\phi_{\theta}x(1-x) + \beta(\psi_{\theta}(1-x)^2 - \psi_{1-\theta}x^2), \quad (7) \quad 24$$

25 where $\phi_{\theta} := 1 - H_{\theta}(\frac{1}{2})$ and $\psi_{\theta} := 1 - H_{\theta}(\hat{p})$. 25
 26

27 Fix $\varepsilon > 0$. Assume that the state switches to $\theta_{t_0} = 1$ at a time where the population 27
 28 has settled on a near-consensus $x_{t_0-1} \simeq 0$. As long as $x_t < \varepsilon$, (7) implies that $x_{t+1} \simeq$ 28
 29 $2\phi_1x_t + \beta\psi_1$ increases at a speed that hinges on β . As $\lambda \rightarrow 0$, the equilibrium value of β 29
 30 converges to zero, and x_{t_0-1} is (with high probability) close to zero, so that the number of 30

1 periods required until $x_t > \varepsilon$ increases. Once $x_t > \varepsilon$, since g_1 is bounded away from the 1
 2 diagonal $y = x$ on the interval $[\varepsilon, 1 - \varepsilon]$, it takes only a finite number of stages, independent 2
 3 of λ , until $x_t > 1 - \varepsilon$. 3

4 The transition dynamics from one consensus to the other thus involves two different 4
 5 phases. In a first phase which grows longer as $\lambda \rightarrow 0$, the old consensus persists despite 5
 6 the state change (inertia): most agents observe a unanimous sample and most likely herd, 6
 7 which slows down society's response. At some point, though, there is enough heterogeneity 7
 8 in the population, and sufficiently many agents draw more balanced samples – enhancing 8
 9 information acquisition – and the society quickly switches to the new consensus in bounded 9
 10 time: there is a *domino* effect whereby the popularity of the new action snowballs. 10

11 Since, as $\lambda \rightarrow 0$, society is almost always in consensus, and information is acquired 11
 12 with vanishing probability, the equilibrium welfare is equal to the fraction of time spent 12
 13 in a correct consensus; in other words, it is directly linked to the duration of the phase of 13
 14 inertia. 14

15 4.3. Equilibrium welfare 15

16 We now focus on equilibrium welfare, and examine how our incomplete learning re- 16
 17 sult extends to larger samples. For simplicity, we assume here that signals are binary with 17
 18 precision $\pi > \frac{1}{2}$. 18
 19 19

20 **THEOREM 3:** *If $\hat{p} = \pi - c$ is high enough, and λ is small enough, there exists an equi-* 20
 21 *librium in which $p_k \in [1 - \hat{p}, \hat{p}]$ for all k : acquiring information is always a best response.* 21
 22 22

23 **PROOF SKETCH:** We look for an equilibrium where $\beta(k) = 1$ for all $k \notin \{0, n\}$ and 23
 24 $\beta := \beta(0) = \beta(n) \in (0, 1)$: agents always acquire information for sure unless they observe 24
 25 a unanimous sample, in which case they randomize between acquiring information and 25
 26 herding on the observed action.¹⁰ We first show that, whenever $n^2\pi(1 - \pi) < 1$, there 26
 27 exists a $\beta > 0$ such that $p_n = 1 - p_0 = \hat{p}$. We next show that, provided \hat{p} is high enough, all 27
 28 other interim beliefs are contained in the interval $[1 - \hat{p}, \hat{p}]$, which completes the proof. 28
 29 29

30 ¹⁰If $c = 0$, $\beta(k)$ are the probabilities of following one's private signal. 30

Using (5) and (6) for such strategies and binary signals, (x_t) obeys $x_t = g_{\theta_t}(x_{t-1})$, with

$$g_{\theta}(x) = \pi_{\theta} + (1 - \beta) \{x^n(1 - \pi_{\theta}) - (1 - x)^n\pi_{\theta}\}, \text{ where } \pi_1 = \pi = 1 - \pi_0.$$

If $\beta = 0$, the analysis of the case $n = 2$ (which holds verbatim) implies that society is eventually trapped in an irreversible consensus. This yields $p_n = p_0 = \frac{1}{2}$, which contradicts $\beta = 0$. If $\beta = \lambda^m$ for an arbitrary m , $p_n \rightarrow 1$ as $\lambda \rightarrow 0$. Indeed, whenever the state switches to $\theta = 1$, (x_t) jumps above $g_1(0) = \beta\pi$ and then increases at a rate of $n\pi > 1$. It thus escapes some fixed neighborhood of 0 in $\ln \frac{1}{\lambda}$ stages and next approaches 1 in boundedly many stages unless the state switches back to $\theta = 0$. For λ small, state changes occur on average every $\frac{1}{\lambda} \gg \ln \frac{1}{\lambda}$ stages; at the steady state, x_t and θ_t are then close to perfectly correlated. This contradicts $\beta > 0$. By a continuity argument, there exists β_{λ} such that $p_n = 1 - p_0 = \hat{p}$.

Such a β_{λ} is an equilibrium iff $p_k \in (1 - \hat{p}, \hat{p})$ for each $k \notin \{0, n\}$. Our methodological contribution consists in providing estimates of such p_k as $\lambda \rightarrow 0$. Bayes rule writes $\frac{p_k}{p_{n-k}} = \frac{I_k}{I_{n-k}}$, with $I_k := \int_0^1 \psi_k(x) dF_1(x)$, where $\psi_k(x) = x^k(1-x)^{n-k}$ and F_{θ} is the invariant cdf conditional on θ — i.e., $F_{\theta}(x) = 2\mu(\theta, [0, x])$ for each x . The argument relies on approximating I_k with $\lambda \sum_{t \in \mathbf{Z}} \psi_k(g_1^t(x))$, where g_1^t is the t -th iterate of g_1 and $(g_1^t(x))_{t \in \mathbf{Z}}$ is a doubly infinite orbit of g_1 . This sum is similar to the time-average of ψ_k during a visit to state $\theta = 1$, hence this approximation is reminiscent of the law of large numbers.

The proof uses the following observation on F_1 . As $\lambda \rightarrow 0$, the probability of a state switch in m periods is vanishingly small, hence the distribution F_1 around $x \in (0, 1)$ coincides with the push-forward measure (given g_1) of F_1 around $g_1^m(x)$ for any fixed m . Specifically, we show that

$$\lim_{\lambda \rightarrow 0} \frac{1}{\lambda} (F_1(y) - F_1(x)) = \lim_{\lambda \rightarrow 0} \frac{1}{\lambda} (F_1(g_1^m(y)) - F_1(g_1^m(x))) \neq 0$$

for any $x, y \in (0, 1)$. This yields $I_k \simeq \lambda \int_z^{g_1(z)} \sum_{t \in \mathbf{Z}} \psi_k(g_1^t(x)) dF_1(x)$ for some z . In the limit $\lambda \rightarrow 0$, this leads to

$$\frac{p_k}{p_{n-k}} \leq \sup_{x \in (0,1)} \frac{\sum_{t \in \mathbf{Z}} \psi_k(\bar{g}_1^t(x))}{\sum_{t \in \mathbf{Z}} \psi_{n-k}(\bar{g}_1^t(x))} \text{ for each } k \notin \{0, n\}, \quad (8)$$

where $\bar{g}_1(\cdot)$ is the function $g_1(\cdot)$ in the specific case where $\beta = 0$. It is straightforward to show that the RHS in (8) is uniformly bounded, which guarantees that $p_k \in (1 - \hat{p}, \hat{p})$ for all $k \notin \{0, n\}$, provided \hat{p} is high enough. *Q.E.D.*

Theorem 3 is reminiscent of the case $n = 2$ because information is acquired following all sample realizations. The underlying reason is similar: if information was not acquired at unanimous samples, society would converge towards a consensus faster than away, making such consensus irreversible, hence uninformative. With $n > 2$, though, the chances of observing a dissenting action in one's sample are higher, which lowers the relative speed of convergence towards the correct consensus. If π is large, though, convergence is still too fast because signals are then more correlated with the true state and, hence, among each other.¹¹ Actions are then less diverse, accelerating convergence.

Equation (8) yields upper and lower bounds on p_k/p_{n-k} (the latter comes from an upper bound on p_{n-k}/p_k). These two bounds are quite close if $n = 3$, allowing for precise estimates of p_1 and p_2 . These equilibrium values are depicted in the right panel of Figure 4, as a function of π . Note that $p_1 > p_2$, which implies that beliefs are non-monotonic in the sample composition.

Some intuition can be found in the left and central panels of Figure 4 (in the case $\pi = 0.99$). The left panel features the functions \bar{g}_θ . Note that $\bar{g}'_0(0)$ is close to 0, hence the population converges quickly towards a consensus on action 0 when $\theta = 0$. The dynamics

¹¹This is why Theorem 3 requires π to be large, unlike Proposition 2 (see the supplementary material for more discussion). In addition, the assumption that c is low helps guarantee that $p_k \in (1 - \hat{p}, \hat{p}_k)$ for $k \notin \{0, n\}$. No such condition is required in Proposition 2 since $p_1 = \frac{1}{2}$.

away from 0 is not nearly as fast when $\theta = 1$. As a result, conditional on x being relatively low, society is more likely to be transitioning away from 0 than towards 0. Hence, the current state is more likely to be $\theta = 1$. This is further illustrated by the central panel, where we plot the logs of (simulated) steady-state, equilibrium densities in the limit $\lambda \rightarrow 0$. We note that this belief reversal $p_1 - p_2$ increases with π , in line with the intuition that convergence to the correct consensus gets relatively faster.




FIGURE 4.—Transition functions, densities, and beliefs for $n = 3$

4.4. The planner's problem

One key question is whether our incomplete learning result is an equilibrium feature or an inescapable feature of the environment. It turns out that a social planner who could dictate any strategy would attain a steady-state welfare of 1 in the persistent limit, so the learning failure is an equilibrium phenomenon. Consider *e.g.* a (symmetric) strategy in which the probability of acquiring information is of order λ for $k = 0$, is 1 for $k = \lfloor \frac{n}{2} \rfloor$ and increases linearly with $k \in \{0, \dots, \frac{n}{2}\}$. Such a strategy ensures that the duration of the phase of inertia is of order $\ln 1/\lambda$, and therefore negligible compared to $1/\lambda$. This ensures that the prevailing consensus is always correct in the limit $\lambda \rightarrow 0$. At the same time, the expected fraction of agents acquiring information vanishes as $\lambda \rightarrow 0$ (see Lévy, Pęski and Vieille (2022) for details).

5. DISCUSSION

In this section, we review the case $\hat{p} = 1$, as well as extensions of our model to asymmetric environments and to a continuum of actions. Other extensions are discussed in the supplementary material.

5.1. The case $\hat{p} = 1$

When $\hat{p} = 1$, that is, signals are free and unbounded, the prevailing consensus is asymptotically correct in the limit $\lambda \rightarrow 0$, as Proposition 3 shows. This result is in line with the fixed-state literature (see, e.g., [Smith and Sørensen \(2000\)](#), [Banerjee and Fudenberg \(2004\)](#)), in contrast to the case $\hat{p} < 1$.

PROPOSITION 3: *Suppose $\hat{p} = 1$ and let $n \geq 1$ be arbitrary. For $\lambda > 0$, let any equilibrium steady state be given, with payoff w^* . Then $\lim_{\lambda \rightarrow 0} w^* = 1$.*

PROOF: As before, we list the actions sampled by a typical agent in some random order $a^{(1)}, \dots, a^{(n)}$. Let $p^{(1)}$ be the interim belief formed on the basis of $a^{(1)}$ only. One strategy consists in replicating $a^{(1)}$. Since $a^{(1)}$ matches yesterday's state with probability w^* , this strategy yields a payoff $u(p^{(1)}) = (1 - \lambda)w^* + \lambda(1 - w^*) \geq w^* - \lambda$. Another strategy consists in ignoring all sampled actions except for $a^{(1)}$ and using one's (free) signal optimally. This strategy yields a payoff $v(p^{(1)})$, which is no larger than the equilibrium payoff w^* . Thus,

$$u(p^{(1)}) \leq v(p^{(1)}) \leq w^* \leq u(p^{(1)}) + \lambda. \quad (9)$$

Since $\hat{p} = 1$, $v(p) > u(p)$ for all $p \in (0, 1)$ and $v(p) = u(p) = 1$ for $p \in \{0, 1\}$. Together with (9), and using the continuity of the functions u and v , this implies $\lim_{\lambda \rightarrow 0} p^{(1)} \in \{0, 1\}$ and $\lim_{\lambda \rightarrow 0} w^* = 1$. *Q.E.D.*

5.2. Asymmetric case

The symmetry assumption made so far is quite convenient, but plays no specific role. Consider a general setup where (a) the probabilities that the state changes from $\theta = 0$ to

1 $\theta = 1$ and from 1 to 0 are λp^* and $\lambda(1 - p^*)$, respectively, so that the invariant probability 1
 2 of state 1 is $p^* \in (0, 1)$, (b) the utility is an arbitrary function $u : \{0, 1\} \times \Theta \rightarrow [0, 1]$, and (c) 2
 3 the (unconditional) distribution of private beliefs is an arbitrary distribution $H \in \Delta([0, 1])$ 3
 4 with expectation $\frac{1}{2}$. 4

5 There exist cutoffs $0 \leq p_0^* \leq p_1^* \leq 1$ such that $v(p) > u(p)$ iff $p_0^* < p < p_1^*$. The results of 5
 6 the paper extend under the assumption that $0 < p_0^* < p^* < p_1^* < 1$. This is a joint assumption 6
 7 on all primitives of the model. The assumption that $p_0^*, p_1^* \in (0, 1)$ states that signals are 7
 8 either bounded or costly (counterpart of Assumption 2). The assumption that $p_0^* < p^* < p_1^*$ 8
 9 ensures that any steady-state equilibrium entails some information acquisition (counterpart 9
 10 of Assumption 1). The detailed statements are in the supplementary material. 10

11 5.3. Continuum of actions 11

12 While we have considered binary actions so far, we provide suggestive evidence that the 12
 13 discontinuity at the limit $\lambda \rightarrow 0$ could arise as well with a richer (infinite) action set. 13

14 For concreteness, assume $A = [0, 1]$ and a square loss utility function $u(a, \theta) = 1 - (a - 14$
 15 $\theta)^2$. Note that, if $c = 0$, one has $\hat{p} = 1$ even with bounded signals because actions are 15
 16 responsive to any extra information; we thus assume $c > 0$ to stick to the assumption $\hat{p} < 1$. 16
 17 17

18 In the fixed-state version where a continuum of new agents sample at least two actions 18
 19 from the past, the distribution of actions converges over time to the correct action (see, *e.g.*, 19
 20 Lee (1993)). With an evolving state, Proposition 4 below shows that the average dispersion 20
 21 of actions in the population vanishes as $\lambda \rightarrow 0$, thereby extending the consensus result of 21
 22 Theorem 2. The proof is in Appendix D. 22

23 PROPOSITION 4: Assume $n \geq 2$ and denote by $a^{(k)}, a^{(l)}$ any two sampled actions. At 23
 24 any equilibrium steady state, one has 24

$$25 \mathbf{E} \left[(a^{(k)} - a^{(l)})^2 \right] \leq \gamma \lambda \quad 25$$

26 where γ is independent of λ and of the equilibrium. 26
 27 27

28 A full-blown analysis of equilibrium behavior and welfare is highly challenging, and 28
 29 beyond the scope of the paper anyway. Without aiming at generality, we discuss here the 29
 30 30

example of *perfect* signals, which yields results consistent with our incomplete learning result. If the state is fixed, all agents acquire information in the first period and convergence to the truth takes exactly one period. When the state is evolving, the population in period t is described by the pair (θ_t, x_t) , where $x_t \in \Delta([0, 1])$ is the distribution of actions in the population. Accordingly, an equilibrium steady state is a distribution $\mu \in \Delta(\Theta \times \Delta([0, 1]))$.

We describe an equilibrium where all agents in period t choose the *same* action $a_t \in [0, 1]$: actions are perfectly correlated, and inferences are independent of the sample size n .

In period $t + 1$, agents then hold the interim belief $f(a_t) := (1 - \lambda)a_t + \lambda(1 - a_t)$. If $f(a_t) \notin [1 - \hat{p}, \hat{p}]$, all agents choose the action $a_{t+1} = f(a_t)$. If instead $f(a_t) \in [1 - \hat{p}, \hat{p}]$, all agents acquire information, learn the state and choose $a_{t+1} = \theta_{t+1} \in \{0, 1\}$. Either way, the consensus is preserved in period $t + 1$. In this equilibrium, agents acquire information periodically: when agents learn that the current state is, say, $\theta_t = 1$, their actions are $a_t = 1$, $a_{t+1} = f(1)$ etc., until they acquire information in period $t + M_\lambda$, where $M_\lambda := \left\lfloor \frac{\ln(2\hat{p} - 1)}{\ln(1 - 2\lambda)} \right\rfloor$. The marginal $\mu_2 \in \Delta(\Delta([0, 1]))$ over action distributions is uniform over the degenerate distributions $\delta_{f^m(\theta)}$ ($0 \leq m < M_\lambda$, $\theta \in \Theta$).

As $\lambda \rightarrow 0$, μ_2 weakly converges to a distribution concentrated over degenerate distributions δ_a , $a \in [0, 1]$. In the persistent limit, actions are perfectly correlated within periods, and the consensus action a has a density: there is no complete learning.¹²

6. RELATION TO THE LITERATURE

Our paper especially connects to five themes in the literature on social learning.¹³

Random sampling. Banerjee and Fudenberg (2004) and Smith and Sørensen (2020) also assume that agents draw a random (finite) sample from a continuum of past actions and identify conditions under which learning is asymptotically complete when the state is fixed. As these papers note, models with a continuum of agents inherently exhibit better aggregative properties than those with a sequence of agents. Indeed, when agents observe a common set of predecessors in one-agent models, *some* histories trigger an incorrect cascade if

¹²One can check that this density is $\frac{1}{2a-1} (1_{[\hat{p}, 1]}(a) - 1_{[0, 1-\hat{p}]}(a))$ up to a normalization constant.

¹³For more complete surveys of this literature, see Smith and Sørensen (2011) and Bikhchandani et al. (2021).

1 signals are bounded. With a continuum of agents, instead, agents update beliefs considering 1
2 *all* possible (mutually exclusive) histories, weighted by their chances (Smith and Sørensen, 2
3 2020). In addition, as soon as agents observe two or more actions, they will rely more on 3
4 their private signals when their sample conveys mixed evidence, which fosters learning. 4
5 With a fixed state, this logic guarantees complete learning even when signals are bounded 5
6 and/or costly (Banerjee and Fudenberg, 2004), and our results thus come in stark contrast. 6

7 *Costly information acquisition.* Our modeling of costly information acquisition relates 7
8 our work to Burguet and Vives (2000) and Ali (2018), who study within one-agent obser- 8
9 vational learning models whether costly private signals preclude (or not) complete learn- 9
10 ing when the state is fixed. Burguet and Vives (2000) endogenize the choice of precision, 10
11 and argue that complete learning arises when information is acquired at beliefs close to 11
12 certainty.¹⁴ Likewise, agents in Ali (2018) can choose from a set of experiments, at an id- 12
13 iosyncratic cost. When costs are bounded away from zero, learning is incomplete as soon 13
14 as signals are bounded. Unlike these papers, we assume a continuum of agents. In this more 14
15 favorable setup, learning is complete when the state is fixed if $n \geq 2$, even if signals are 15
16 costly (see our extension of Banerjee and Fudenberg (2004) to costly signals in the supple- 16
17 mentary material), in contrast to our main results. The example we analyze in Section 5.3 17
18 of a model with a continuum of (responsive) actions reinforces our message: costly signals 18
19 do not preclude complete learning with a fixed state, but concur to incomplete learning with 19
20 changing states. 20

21 *Changing states.* Our paper also relates to a stream of papers on social learning in a 21
22 changing world. Moscarini, Ottaviani and Smith (1998) show that cascades must end in 22
23 finite time, but arise for sure when the state is persistent enough. Based on their setup, 23
24 Huang (2022) shows that in the long run the frequency of action changes is higher than that 24
25 of state changes. In a different line of research, Ellison and Fudenberg (1995), Acemoglu, 25
26 Nedic and Ozdaglar (2008) and Frongillo, Schoenebeck and Tamuz (2011) consider non- 26
27 Bayesian models in which agents follow specific heuristics. 27

28
29
30 ¹⁴This is in line with our discussion of endogenous information structures, see the (online) Section H. 30

1 *Stationary analyses of social learning.* [Dasaratha, Golub and Hak \(2023\)](#) and [Kabos and](#) 1
2 [Meyer \(2021\)](#) also develop stationary analyses of social learning. [Dasaratha, Golub and](#) 2
3 [Hak](#) consider a Gaussian environment where agents in a network learn from their neigh- 3
4 bors. They show that learning is improved when agents have heterogeneous neighbors who 4
5 have access to signals of different precision. While we rule out such heterogeneity, our 5
6 analysis also highlights the adverse welfare impact of an excessive correlation of actions 6
7 between (symmetric) agents. [Kabos and Meyer](#) consider a Markovian environment where 7
8 past actions may be misrecorded, and investigate whether agents put too much or too little 8
9 weight on their private information, while we focus on providing bounds on equilibrium 9
10 welfare. 10

11 *Rate of learning and the Grossman-Stiglitz paradox.* Our results echo the well-known 11
12 Grossman-Stiglitz paradox ([Grossman and Stiglitz, 1980](#)) according to which agents would 12
13 ignore their individual signals if information was fully aggregated, precluding information 13
14 aggregation in the first place. In a fixed-state world where asymptotic learning is guaran- 14
15 teed, such a logic imposes that learning be necessarily slow, as shown in [Vives \(1993\)](#). 15
16 In a social learning context, [Harel et al. \(2021\)](#) also establish slow social learning even if 16
17 agents observe several actions, because the correlation in the agents' actions arising from 17
18 social learning reduces the amount of information these actions reveal about the state (see 18
19 also [Huang, Strack and Tamuz \(2024\)](#) on slow convergence). While a high correlation re- 19
20 duces the speed of learning with a fixed state, it lowers responsiveness, hence steady-state 20
21 welfare, in our changing state environment. 21

22 23 24 25 26 27 28 29 30

7. CONCLUSION

25 We consider a general model of social learning with binary actions and states in which 25
26 states change over time, information is possibly costly, and agents draw finite samples 26
27 of past actions. We show that, under a wide range of situations, the possibility that the 27
28 state changes drastically limits the extent of social learning at the steady state, in crisp 28
29 contrast to what would happen in analogous fixed-state environments. Beyond this insight, 29
30 the methods we develop could pave the way to address interesting questions on how a 30

1 planner would optimally design the learning environment (sampling procedures, feedback 1
 2 given to players,...) to foster the welfare gains from social learning. 2

3
 4 REFERENCES 4

5 **Acemoglu, Daron, Angelia Nedic, and Asuman Ozdaglar.** 2008. “Convergence of rule-of-thumb learning rules
 6 in social networks.” 1714–1720, IEEE. [2, 24]

7 **Ali, S Nageeb.** 2018. “Herding with costly information.” *Journal of Economic Theory*, 175: 713–729. [24]

8 **Banerjee, Abhijit, and Drew Fudenberg.** 2004. “Word-of-mouth learning.” *Games and economic behavior*,
 9 46(1): 1–22. [3, 12, 21, 23, 24]

10 **Bhattacharya, Rabi, and Mukul Majumdar.** 2007. *Random dynamical systems: theory and applications*. Cam-
 11 bridge University Press. [13]

12 **Bikhchandani, Sushil, David Hirshleifer, Omer Tamuz, and Ivo Welch.** 2021. “Information cascades and
 13 social learning.” National Bureau of Economic Research. [1, 23]

14 **Burguet, Roberto, and Xavier Vives.** 2000. “Social learning and costly information acquisition.” *Economic*
 15 *theory*, 15(1): 185–205. [24]

16 **Dasaratha, Krishna, Benjamin Golub, and Nir Hak.** 2023. “Learning from neighbours about a changing state.”
 17 *Review of Economic Studies*, 90(5): 2326–2369. [2, 25]

18 **Ellison, Glenn, and Drew Fudenberg.** 1995. “Word-of-mouth communication and social learning.” *The Quar-*
 19 *terly Journal of Economics*, 110(1): 93–125. [24]

20 **Frongillo, Rafael M, Grant Schoenebeck, and Omer Tamuz.** 2011. “Social learning in a changing world.”
 21 146–157, Springer. [2, 24]

22 **Grossman, Sanford J, and Joseph E Stiglitz.** 1980. “On the impossibility of informationally efficient markets.”
 23 *The American economic review*, 70(3): 393–408. [25]

24 **Harel, Matan, Elchanan Mossel, Philipp Strack, and Omer Tamuz.** 2021. “Rational groupthink.” *The Quar-*
 25 *terly Journal of Economics*, 136(1): 621–668. [25]

26 **Huang, Wanying.** 2022. “The Emergence of Fads in a Changing World.” *arXiv preprint arXiv:2208.14570*. [2,
 27 24]

28 **Huang, Wanying, Philipp Strack, and Omer Tamuz.** 2024. “Learning in repeated interactions on networks.”
 29 *Econometrica*, 92(1): 1–27. [25]

30 **Kabos, Eszter, and Margaret Meyer.** 2021. “A Welfare Analysis of a Steady-State Model of Observational
 Learning.” working paper. [25]

Lee, In Ho. 1993. “On the convergence of informational cascades.” *Journal of Economic theory*, 61(2): 395–411.
 [22]

Lévy, Raphaël, Marcin Peški, and Nicolas Vieille. 2022. “Stationary social learning in a changing environment.”
 arXiv:2201.02122. [20]

- 1 **Moscarini, Giuseppe, Marco Ottaviani, and Lones Smith.** 1998. “Social learning in a changing world.” *Eco-* 1
 2 *nomical Theory*, 11(3): 657–665. [2, 24] 2
- 3 **Smith, Lones, and Peter Norman Sørensen.** 2011. “Observational learning.” *The New Palgrave Dictionary of* 3
 4 *Economics Online Edition*, 29–52. [23] 4
- 5 **Smith, Lones, and Peter Norman Sørensen.** 2020. “Rational Social Learning with Random Sampling.” working 5
 6 paper. [23, 24] 6
- 7 **Smith, Lones, and Peter Sørensen.** 2000. “Pathological outcomes of observational learning.” *Econometrica*, 68 7
 8 (2): 371–398. [21] 8
- 9 **Solomyak, Boris.** 1995. “On the random series $\sum \pm \lambda^n$ (an Erdos problem).” *Annals of Mathematics*, 611–625. 9
 10 [13] 10
- 11 **Vives, Xavier.** 1993. “How fast do rational agents learn?” *The Review of Economic Studies*, 60(2): 329–347. [25] 11

APPENDIX A: THE CASE $n = 2$: PROPOSITION 2

13 We argue by contradiction and assume that $p_2 > \hat{p}$ for some equilibrium steady state. 13
 14 Since agents who observe a balanced sample $k = 1$ hold the belief $p_1 = \frac{1}{2}$ and acquire 14
 15 information, the fraction of agents choosing action 1 in period $t + 1$ reads 15

$$x_{t+1} = x_t^2 + 2x_t(1 - x_t)\phi_{\theta_{t+1}},$$

16 where $\phi_\theta = 1 - H_\theta(\frac{1}{2})$ is the probability of playing action 1 in state θ when holding an 16
 17 interim belief $\frac{1}{2}$. Note that $\phi_1 = 1 - \phi_0 > \frac{1}{2}$. 17

18 **LEMMA A.1:** *The sequence (x_t) converges a.s., with $\lim_{t \rightarrow +\infty} x_t \in \{0, 1\}$.* 18

19 **PROOF:** Choose $\tilde{\phi}_\theta > \phi_\theta$ such that $\tilde{\phi}_0\tilde{\phi}_1 < \frac{1}{4}$, and $\varepsilon_0 > 0$ such that $2\phi_\theta + \varepsilon_0 < 2\tilde{\phi}_\theta$ for 19
 20 each θ . Note that $g_\theta(x) \leq 2\tilde{\phi}_\theta x$ for $x < \varepsilon_0$. 20

21 Let $\varepsilon < \varepsilon_0$ be arbitrary. We define two interlacing sequences $(\tau_m^{\text{in}})_m$ and $(\tau_m^{\text{out}})_m$ of 21
 22 stopping times. We first set 22

$$\tau_1^{\text{out}} := \inf\{t \geq 0 : x_t < \varepsilon \text{ and } \theta_t = 0, \text{ or } x_t > 1 - \varepsilon \text{ and } \theta_t = 1\},$$

23 and $\tau_1^{\text{in}} := \inf\{t \geq \tau_1^{\text{out}} : x_t \in [\varepsilon, 1 - \varepsilon]\}$, with $\inf \emptyset = +\infty$. 23
 24 24
 25 25
 26 26
 27 27
 28 28
 29 29
 30 30

These are the first exit and entry times in $[\varepsilon, 1 - \varepsilon]$.¹⁵ For $m \geq 1$, we set

$$\tau_{m+1}^{\text{out}} := \inf\{t \geq \tau_m^{\text{in}} : x_t < \varepsilon \text{ and } \theta_t = 0, \text{ or } x_t > 1 - \varepsilon \text{ and } \theta_t = 1\},$$

and $\tau_{m+1}^{\text{in}} := \inf\{t \geq \tau_{m+1}^{\text{out}} : x_t \in [\varepsilon, 1 - \varepsilon]\}$.

Below, we show that (x_t) cannot remain indefinitely in the interval $[\varepsilon, 1 - \varepsilon]$.

CLAIM 4: *One has $\mathbf{P}(\tau_{m+1}^{\text{out}} < +\infty \mid \tau_m^{\text{in}} < +\infty) = 1$ for each m .*

PROOF: For $x_t = x \in [\varepsilon, 1 - \varepsilon]$, one has

$$g_1(x) - x = x - g_0(x) = (2\phi_1 - 1)(x - x^2) \geq (2\phi_1 - 1)\varepsilon(1 - \varepsilon) \quad (\text{A.1})$$

and therefore the difference $x_{t+1} - x_t$ is bounded away from zero. With $N := \lceil \frac{1}{\varepsilon((1 - \varepsilon)2\phi_1 - 1)} \rceil$ and using (A.1), it follows that $x_t \in [\varepsilon, 1 - \varepsilon]$ implies $x_{t+N} \notin [\varepsilon, 1 - \varepsilon]$ as soon as $\theta_{t+1} = \dots = \theta_{t+N}$, which has probability $(1 - \lambda)^N$. This implies

$$\mathbf{P}(\tau_{m+1}^{\text{out}} \leq t + N \mid \tau_m^{\text{in}} \leq t < \tau_{m+1}^{\text{out}}) \geq (1 - \lambda)^N,$$

and therefore,

$$\mathbf{P}(\tau_{m+1}^{\text{out}} \geq t + jN \mid \tau_m^{\text{in}} \leq t < \tau_{m+1}^{\text{out}}) \leq \left(1 - (1 - \lambda)^N\right)^j$$

for each j . The result follows when $j \rightarrow +\infty$. *Q.E.D.*

We show that the probability that x_t ever re-enters the interval $[\varepsilon, 1 - \varepsilon]$ after it exits from it, is bounded away from 1. In the next statement, $(\mathcal{H}_t)_t$ is the natural filtration of $(\theta_t, x_t)_t$ and $\mathcal{H}_{\tau_m^{\text{out}}}$ is the stopped filtration at time τ_m^{out} .

CLAIM 5: *There exists $a > 0$ such that for each m , $\mathbf{P}(\tau_m^{\text{in}} = +\infty \mid \mathcal{H}_{\tau_m^{\text{out}}}) \geq a$, w.p. 1 on the event $\tau_m^{\text{out}} < +\infty$.*

PROOF: Consider the event $\tau_m^{\text{out}} = t$. We assume for concreteness that $x_t < \varepsilon$ and $\theta_t = 0$. By the Markov property, we may assume w.l.o.g. that $t = 0$ and $m = 1$.

¹⁵Except for the extra condition on the exit state in the definition of τ_1^{out} , which is for convenience.

We define an auxiliary sequence (W_t) of random variables by $W_0 = 0$ and $W_{t+1} = W_t + \ln 2\tilde{\phi}_{\theta_{t+1}}$ for $t \geq 1$. Since $x_{t+1} \leq 2x_t\tilde{\phi}_{\theta_{t+1}}$, one has $W_t \geq \ln x_t - \ln x_0$ for each t . This implies that $\tau_1^{\text{in}} \geq \inf\{t \geq 1 : W_t \geq 0\}$, and $\mathbf{P}(\tau_1^{\text{in}} < +\infty) \leq \mathbf{P}(\sup_{t \geq 1} W_t \geq 0)$. Introduce the successive state changes $\psi_0 = 0$ and $\psi_{m+1} := \inf\{t > \psi_m : \theta_t \neq \theta_{t-1}\}$. Assuming $\theta_0 = 0$, $\theta_t = 1$ whenever $\psi_{2m-1} \leq t < \psi_{2m}$ for some m , and $\theta_t = 0$ otherwise.¹⁶

For $i \in \mathbf{N}$, set $X_i := W_{\psi_{2i+2}} - W_{\psi_{2i}}$. Observe that $W_t - W_{t-1} > 0$ iff $\theta_t = 1$, hence

$$\sup_t W_t \geq 0 \Leftrightarrow \sup_i (X_0 + \cdots + X_i) \geq 0.$$

The r.v.'s (X_i) are *i.i.d.* with $\mathbf{E}[X_1] = \frac{1}{\lambda} (\ln 2\tilde{\phi}_1 + \ln 2\tilde{\phi}_0) < 0$. The sequence $(X_0 + \cdots + X_j)_j$ is therefore a simple random walk with negative drift, which implies

$$\mathbf{P}\left(\sup_j (X_1 + \cdots + X_j) \geq 0\right) \leq 1 - a \text{ for some } a > 0.$$

Q.E.D.

Claims 4 and 5 yield $\mathbf{P}(\tau_m^{\text{out}} < +\infty \text{ and } \tau_{m+1}^{\text{in}} = +\infty \text{ for some } m) = 1$. Hence there is a.s. finite random time T_0 such that either $x_t < \varepsilon$ for all $t \geq T_0$, or $x_t > 1 - \varepsilon$ for all $t \geq T_0$. *Q.E.D.*

LEMMA A.2: *The only symmetric invariant measure for (θ_t, x_t) is uniform over $\Theta \times \{0, 1\}$.*

PROOF: By Lemma A.1, any invariant measure is concentrated on $\Theta \times \{0, 1\}$.¹⁷ Since the sets $\{x = 0\}$ and $\{x = 1\}$ are absorbing for (θ_t, x_t) , one has for $a \in \{0, 1\}$

$$\begin{aligned} \mu(0, a) &= \mathbf{P}((\theta_{t+1}, x_{t+1}) = (0, a)) \\ &= (1 - \lambda)\mathbf{P}((\theta_t, x_t) = (0, a)) + \lambda\mathbf{P}((\theta_t, x_t) = (1, a)) \\ &= (1 - \lambda)\mu(0, a) + \lambda\mu(1, a). \end{aligned}$$

¹⁶If $\theta_0 = 1$, odd and even phases should be switched.

¹⁷Indeed, by the invariance property, $\mu(\Theta \times [\varepsilon, 1 - \varepsilon]) = \mathbf{P}(x_t \in [\varepsilon, 1 - \varepsilon])$ for each t . By Lemma A.1, the RHS converges to zero as $t \rightarrow +\infty$ for each $\varepsilon > 0$.

Hence $\mu(0, a) = \mu(1, a)$: μ is a product distribution. Since μ is symmetric, it is uniform.

Q.E.D.

The result follows. By Lemma A.2, $p_2 = \frac{1}{2}$ – a contradiction.

APPENDIX B: EQUILIBRIUM EXISTENCE: THEOREM 1

We prove existence using a fixed-point argument. Define:

- M to be the set of distributions $\mu \in \Delta(\Theta \times [0, 1])$ that are invariant when permuting the two states.
- B to be the set of $b = (b_s) \in [0, 1]^{\{0, \dots, n\}}$ such that $b_s = 1 - b_{n-s}$ for each s , with the interpretation that b_s is the interim belief that $\theta = 1$ when sampling s .
- Φ to be the set of $\phi = (\phi_{\theta, s}) \in [0, 1]^{\Theta \times \{0, \dots, n\}}$ such that $\phi_{0, s} = 1 - \phi_{1, n-s}$ for each s , with the interpretation that $\phi_{\theta, s}$ is the overall probability of playing action 1, when in state θ and sampling s .

M is compact metric when endowed with the topology of weak convergence, and $\Sigma := M \times B \times \Phi$ is convex compact with the product topology. We define a set-valued map $\Psi : \Sigma \rightarrow \Sigma$ by $\Psi(\mu, b, \phi) := \psi_1(\phi) \times \psi_2(\mu) \times \psi_3(b)$, where ψ_1, ψ_2, ψ_3 are defined below.

Definition and properties of ψ_1

Let $\phi \in \Phi$ be given. It induces a (symmetric) Markov chain (θ_t, x_t) on $\Omega := \Theta \times [0, 1]$, with (x_t) obeying the recursive equation

$$x_{t+1} = (1 - \rho)x_t + \rho \sum_{s=0}^n \text{bin}_{n, x_t}(s) \phi_{\theta_t, s}, \text{ with } \text{bin}_{n, x}(s) := \binom{n}{s} x^s (1 - x)^{n-s}.$$

We set $\psi_1(\phi) := \{\mu \in M : \mu \text{ is invariant for } (\theta_t, x_t)\}$.

LEMMA 2: *The map ψ_1 is uhc, with non-empty convex values.*

PROOF: The proof is standard and only sketched. For given $\phi \in \Phi$, denote $P_\phi(\omega, d\omega')$ the one-step transition probability of (θ_t, x_t) . Note that the support of $P_\phi(\omega, d\omega')$ is a two-point set, for each $\omega \in \Omega$.

For fixed ϕ and $f \in C(\Omega)$, the map $Tf(\omega) = \int_{\omega} f(\omega') P_{\phi}(\omega, d\omega')$ is continuous over Ω . This implies that $\mu \in M \mapsto \mu P_{\phi}$ is continuous in the weak-* topology, and thus has a fixed point by Tychonov Theorem. Thus, $\psi_1(\phi) \neq \emptyset$.

The same argument shows that $(\phi, \mu) \in \Phi \times M \mapsto \mu P_{\phi}$ is continuous as well and linear in μ . This completes the proof. *Q.E.D.*

Definition and properties of ψ_2

Fix $\mu \in M$. The probability of sampling $s \in \{0, 1, \dots, n\}$ in state θ is $P_{\theta}(s | \mu) := \int_{[0,1]} \text{bin}_{n,x}(s) \mu_{\theta}(dx)$, where μ_{θ} is the conditional distribution of x given θ .¹⁸ Denote by $\psi_2^s(\mu)$ the set of all beliefs $b_s \in [0, 1]$ that are consistent with Bayesian updating, that is,

$$\psi_2^s(\mu) := \{b_s \in [0, 1] : P_1(s | \mu)(1 - b_s) = P_0(s | \mu)b_s\},$$

and set

$$\psi_2(\mu) := \left\{ b \in \prod_s \psi_2^s(\mu) : b_s = 1 - b_{n-s} \text{ for each } s \right\}.$$

The proof of Lemma 3 below is straightforward.

LEMMA 3: *The map $\psi_2 : M \rightarrow B$ is uhc, with non-empty convex values.*

Definition and properties of ψ_3

Fix $b \in B$. For given s, θ , we define $\psi_3^{s,\theta}(b_s)$ as the probability of playing action 1 when holding the belief b_s in state θ . Formally,

- $\psi_3^{s,\theta}(b_s) = \{1\}$ if $b_s > \hat{p}$ and $\psi_3^{s,\theta}(b_s) = \{0\}$ if $b_s < 1 - \hat{p}$.
- $\psi_3^{s,\theta}(b_s) = [1 - H_{\theta}(1 - b_s), 1 - H_{\theta}(1 - b_s) -]$ if $b_s \in (1 - \hat{p}, \hat{p})$.
- $\psi_3^{s,\theta}(\hat{p}) := [1 - H_{\theta}(1 - b_s), 1]$ and $\psi_3^{s,\theta}(1 - \hat{p}) := [0, 1 - H_{\theta}(1 - b_s) -]$.

We next set

$$\psi_3(b) := \{\phi \in \Phi : \phi_{\theta,s} \in \psi_3^{\theta,s}(b_s) \text{ for each } (\theta, s) \text{ and } \phi_{0,s} = 1 - \phi_{1,n-s} \text{ for each } s\}.$$

¹⁸By the symmetry assumption, θ is uniformly distributed under μ , so the conditional distribution is well-defined.

The proof of Lemma 4 is straightforward.

LEMMA 4: *The map $\psi_3 : B \rightarrow \Phi$ is upper hemi-continuous with non-empty convex values.*

By Kakutani-Fan-Glicksberg Theorem, Ψ has a fixed point. By construction, the fixed points of Ψ are the symmetric equilibrium steady states. We conclude by proving that there is an equilibrium (σ, μ) such that μ has no atom at 0 and 1.¹⁹

LEMMA 5: *Let (σ, μ) be an equilibrium. There exists an invariant distribution μ' for σ with no atom at 0 nor 1, such that (σ, μ') is an equilibrium.*

PROOF: We start with two observations. Let $g_\theta : [0, 1] \rightarrow [0, 1]$ describe the evolution of x induced by σ . If $g_\theta(x) = 0$ for some $x \in (0, 1)$ and $\theta \in \Theta$, then this holds for all $(x, \theta) \in [0, 1] \times \Theta$ – contradicting the symmetry assumption. In addition, $g_\theta(0) = 0$ if and only if $\beta(0) = 0$ (agents with a unanimous sample do not acquire information).

Assume that μ has an atom at 0. Since $g_\theta(x) > 0$ for all $x \in (0, 1)$, this implies that $g_\theta(0) = 0$ and therefore that $\beta(0) = 0$. Write $\mu = \rho_* \mu_* + (1 - \rho_*) \mu'$, where μ_* is the uniform distribution over $\Theta \times \{0, 1\}$ and μ' has no atom at 0 nor 1. Since $\beta(0) = \beta(n) = 0$, μ_* is invariant for σ and therefore, μ' as well. Since samples are a.s. unanimous under μ_* , the interim beliefs p_k for $k \neq 0$ and $k \neq n$ are the same, whether they are computed using μ or μ' . In addition, p_n is higher when computed with μ than when computed with μ_* and is therefore even higher when computed using μ' , and a symmetric property holds for p_0 . This implies that (σ, μ') is an equilibrium, as desired.

¹⁹We note that whenever there is an equilibrium (σ, μ) with welfare $w^* > \hat{p}$, there is an additional, atomic one with welfare \hat{p} provided λ is small. Indeed, if $w^* > \hat{p}$ and λ is small, agents do not acquire information with a unanimous sample. Consider now the non-informative, uniform distribution μ_* over $\Theta \times \{0, 1\}$. It is invariant for σ , and therefore any convex distribution $\mu_r := r\mu_* + (1 - r)\mu$ as well. For $k \notin \{0, n\}$, interim beliefs $p_k(r)$ inferred from μ_r are constant in r , while $p_n(r)$ is decreasing in r . Thus, (σ, μ_r) is also an equilibrium, provided r is s.t. $p_n(r) \geq \hat{p}$, and the welfare spans the interval $[\hat{p}, w^*]$. We view this source of equilibrium multiplicity as spurious. Indeed, such equilibria are ruled out when requiring that equilibrium distributions μ have no atom at 0 and 1 (such equilibria exist) with no impact on our results.

Q.E.D.

APPENDIX C: NO SOCIAL LEARNING: THEOREM 3

C.1. Notation and Preliminaries

Slightly enriching the notation from the text, let us set

$$g_\theta^\beta(x) = \pi_\theta + (1 - \beta) \{x^n (1 - \pi_\theta) - (1 - x)^n \pi_\theta\} \quad (\text{with } \pi_1 = \pi = 1 - \pi_0).$$

We note that g_θ^β is an increasing bijection from $[0, 1]$ to $[\pi_\theta \beta, \pi_\theta + (1 - \beta)(1 - \pi_\theta)]$, and that there is a unique \bar{x}_θ^β such that $g_\theta^\beta(\bar{x}_\theta^\beta) = \bar{x}_\theta^\beta$. In addition, g_1^β is concave on $[0, 1]$, $(g_1^\beta)'$ is convex on $[0, 1]$ and $g_1^\beta(x) - x$ is decreasing on $[\frac{1}{2}, 1]$.

We denote by $h_\theta^\beta : [g_\theta(\bar{x}_0^\beta), g_\theta(\bar{x}_1^\beta)] \rightarrow [\bar{x}_0^\beta, \bar{x}_1^\beta]$ the inverse of g_θ^β on $[g_\theta(\bar{x}_0^\beta), g_\theta(\bar{x}_1^\beta)]$. We will view h_θ^β as a function defined on $[\bar{x}_0^\beta, \bar{x}_1^\beta]$ by setting

$$h_0^\beta(x) = \bar{x}_1^\beta \text{ for } x > g_0^\beta(\bar{x}_1^\beta) \text{ and } h_1^\beta(x) = \bar{x}_0^\beta \text{ for } x < g_1^\beta(\bar{x}_0^\beta).$$

Given λ , any invariant measure $\mu_{\lambda, \beta}$ for the strategy β is concentrated on $\Theta \times (\bar{x}_0^\beta, \bar{x}_1^\beta)$. We denote by $F_\theta^{\lambda, \beta}(x) := 2\mu_{\lambda, \beta}(\{\theta\} \times [0, x])$ the cdf of the population state given θ .

To avoid clumsy notation, we henceforth omit the superscripts β and λ and simply write $\bar{x}_\theta, \mu, F_\theta, g_\theta, h_\theta$ and p_k . The m -th iterate of g_θ is denoted g_θ^m .

Given a (symmetric) invariant measure μ for β , time invariance implies that

$$F_\theta(x) = (1 - \lambda)F_\theta(h_\theta(x)) + \lambda F_{1-\theta}(h_{1-\theta}(x)) \text{ for each } \theta \in \Theta \text{ and } x \in [\bar{x}_0, \bar{x}_1]. \quad (\text{C.1})$$

LEMMA C.1: For each $x \in [\bar{x}_0, \bar{x}_1]$ and $\theta \in \Theta$, one has

$$|F_\theta(x) - F_\theta(h_\theta(x))| \leq \lambda \text{ and } |F_\theta(x) - F_\theta(g_\theta(x))| \leq \lambda$$

PROOF: The first claim follows from (C.1), the second when applying the first to $g_\theta(x)$.

Q.E.D.

LEMMA C.2: Let $\varepsilon > 0$ be given. There exists k such that for each β small enough, the following holds. For each $x \geq \varepsilon$, $y \in [x, g_1(x)]$, $\lambda > 0$ and $m \in \mathbf{N}$, one has

$$\left| \frac{1}{\lambda} (F_1(g_1^m(y)) - F_1(g_1^m(x))) - \frac{1}{\lambda} (F_1(y) - F_1(x)) \right| \leq mk\lambda.$$

In addition, for each $x, y \in (0, 1)$, one has

$$\lim_{\lambda \rightarrow 0} (F_1(y) - F_1(x)) = 0.$$

PROOF: Let $\varepsilon > 0$, and assume that $n\pi\beta \leq \frac{1}{2}\varepsilon$. There exists k s.t. $g_0^k(g_1(x)) \leq x$, hence $h_0(g_1(x)) \leq h_0^{k+1}(x)$, for each $x \geq \varepsilon$. By Lemma C.1,

$$F_0(h_0(y)) \leq F_0(h_0(g_1(x))) \leq F_0(h_0^{k+1}(x)) \leq F_0(h_0(x)) + k\lambda$$

hence

$$F_0(h_0(y)) - F_0(h_0(x)) \leq k\lambda. \quad (\text{C.2})$$

Applying (C.1) repeatedly, we obtain:

$$\begin{aligned} & F_1(g_1^m(y)) - F_1(g_1^m(x)) \\ &= (1 - \lambda) (F_1(g_1^{m-1}(y)) - F_1(g_1^{m-1}(x))) + \lambda (F_0(h_0(g_1^m(y))) - F_0(h_0(g_1^m(x)))) \\ &= (1 - \lambda)^m (F_1(y) - F_1(x)) + \lambda \sum_{i=1}^m (1 - \lambda)^{i-1} (F_0(h_0(g_1^i(y))) - F_0(h_0(g_1^i(x)))) \\ &\leq F_1(y) - F_1(x) + mk\lambda^2, \end{aligned}$$

where the last inequality follows from (C.2), since $g_1^i(y) \in [g_1^i(x), g_1(g_1^i(x))]$ for each i .

Under the same conditions on β , for given x and y there exists m such that $y \leq g_1^m(x)$. Using Lemma C.1, one has $F_1(x) \leq F_1(y) \leq F_1(x) + k\lambda$ for each $\lambda > 0$. *Q.E.D.*

C.2. The choice of β

Given λ small, we prove here the existence of a strategy β_λ such that agents with a unanimous sample are indifferent whether to acquire information.²⁰

²⁰The existence result requires no assumption on \hat{p} , beyond $\hat{p} \in (\frac{1}{2}, 1)$, but the value of β_λ of course depends on \hat{p} .

1 PROPOSITION C.1: For any $m > 0$ and λ small enough, there exists $\beta_\lambda \leq \lambda^m$, and an
 2 invariant measure for β_λ , such that $p_n = 1 - p_0 = \hat{p}$.

3
 4 The result follows from Lemmas C.3, C.4 and C.6, and from the fact that the set of
 5 symmetric invariant measures is convex-valued and upper hemi-continuous as a function
 6 of $\beta \in [0, 1]$.

7 LEMMA C.3: Let $\beta = 0$. Then $p_n = p_0 = \frac{1}{2}$, for each λ .

8
 9 PROOF: Since $n^2\pi(1 - \pi) < 1$, the statements of Lemmas 2 and 3, and Claims 4 and
 10 5 from Appendix A remain true verbatim. The only (symmetric) invariant measure is the
 11 uniform distribution over $\Theta \times \{0, 1\}$. Q.E.D.

12
 13 LEMMA C.4: Let $m > 0$ be given. For $\lambda > 0$, set $\beta = \lambda^m$. Then $\lim_{\lambda \rightarrow 0} p_n = 1$.

14
 15 PROOF: Observe that $g_1(\bar{x}_0) \geq \beta\pi = \lambda^m\pi$. On the other hand, since $g_1'(0) = (1 -$
 16 $\beta)n\pi > 1$, there exists $\varepsilon > 0$ and $a_1 > 1$ such that $g_1(x) \geq a_1x$ for every $x \in [\bar{x}_0, \varepsilon]$ and
 17 λ small enough. Hence there exists some M independent of λ such that $g_1^i(\bar{x}_0) > \varepsilon$ for
 18 each $i \geq M \ln \frac{1}{\lambda}$. Using Lemma C.1, it follows that

$$19 \quad F_1(\varepsilon) \leq F_1(\bar{x}_0) + \sum_{i=0}^{\lfloor M \ln \frac{1}{\lambda} \rfloor} F_1(g_1^{i+1}(\bar{x}_0)) - F_1(g_1^i(\bar{x}_0)) \leq M\lambda \ln \frac{1}{\lambda}$$

20
 21 hence $\lim_{\lambda \rightarrow 0} F_1(\varepsilon) = 0$. Using Lemma C.2, this implies $\lim_{\lambda \rightarrow 0} F_1(x) = 0$ for each $x < 1$
 22 and by symmetry, $\lim_{\lambda \rightarrow 0} F_0(x) = 1$ for each $x > 0$. The result follows. Q.E.D.

23 C.3. Estimates on F

24
 25 From now on, we set $\beta = \beta_\lambda$. We here derive further estimates on the invariant measure.

26
 27 LEMMA C.5: For each $x \in (0, 1)$, one has

$$28 \quad \lim_{\lambda \rightarrow 0} \frac{1}{\lambda} (F_1(g_1(x)) - F_1(x)) = 2\hat{p} - 1.$$

29
 30

PROOF: Let $\varepsilon > 0$ be given. By Lemma C.1, one has

$$\lim_{\lambda \rightarrow 0} \left(\int_0^1 (1-x)^n dF_\theta(x) - F_\theta(\varepsilon) \right) = 0.$$

Since $\frac{\int_0^1 (1-x)^n dF_1(x)}{\int_0^1 (1-x)^n dF_0(x)} = \frac{p_0}{1-p_0} = \frac{1-\hat{p}}{\hat{p}}$, this implies that $F_\theta(\varepsilon)$ is bounded away from zero as $\lambda \rightarrow 0$, with $\lim_{\lambda \rightarrow 0} \frac{F_1(\varepsilon)}{F_0(\varepsilon)} = \frac{1-\hat{p}}{\hat{p}}$.

On the other hand, symmetry and Lemma C.1 imply that $\lim_{\lambda \rightarrow 0} F_1(\varepsilon) = 1 - \lim_{\lambda \rightarrow 0} F_0(\varepsilon)$. Thus, $\lim_{\lambda \rightarrow 0} F_1(\varepsilon) = \lim_{\lambda \rightarrow 0} F_1(x) = 1 - \hat{p}$ for each $x \in (0, 1)$, and $\lim_{\lambda \rightarrow 0} F_0(\varepsilon) = \lim_{\lambda \rightarrow 0} F_0(x) = \hat{p}$ for each $x \in (0, 1)$. It follows that

$$\lim_{\lambda \rightarrow 0} \frac{1}{\lambda} (F_1(g_1(x)) - F_1(x)) = \lim_{\lambda \rightarrow 0} (F_0(h_0(x)) - F_1(h_1(x))) = 2\hat{p} - 1.$$

Q.E.D.

LEMMA C.6: For each $\varepsilon > 0$ and θ , one has $\limsup_{\lambda \rightarrow 0} (\lambda\varepsilon + \lambda^2)^{-1} \int_0^\varepsilon x dF_\theta(x) < \infty$.

PROOF: Fix $\varepsilon > 0$, and assume for concreteness $\theta = 0$. The other case follows a similar logic. Note first that $\int_0^\varepsilon x dF_0(x) \leq \beta_\lambda \lambda^{-1} + \int_{\beta_\lambda \lambda^{-1}}^\varepsilon x dF_0(x)$. Since $\beta_\lambda \leq \lambda^3$, the first integral is at most λ^2 for λ small.

On the other hand, note that $g_0(x) \leq \bar{x}_0 + (\max_{[\bar{x}_0, x]} g'_0)(x - \bar{x}_0)$ for each $x \in [\bar{x}_0, \bar{x}_1]$.²¹ Since $g'(0) = (1 - \beta_\lambda)n(1 - \pi) < 1$, and provided ε is small enough, there exists some $a_0 < 1$ independent of λ such that $g_0(x) < a_0 x$ for each $x \in [\bar{x}_0, \varepsilon]$ and λ small. Using Lemma C.1, one therefore has

$$\begin{aligned} \int_{\beta_\lambda \lambda^{-1}}^\varepsilon x dF_0(x) &\leq \sum_{m: g_0^m(\varepsilon) > \beta_\lambda \lambda^{-1}} g_0^m(\varepsilon) [F_0(g_0^m(\varepsilon)) - F_0(g_0^{m+1}(\varepsilon))] \\ &\leq \lambda \sum_{m: g_0^m(\varepsilon) > \beta_\lambda \lambda^{-1}} g_0^m(\varepsilon) \leq \lambda \sum_m a_0^m \varepsilon \leq \frac{1}{1-a_0} \lambda \varepsilon. \end{aligned}$$

Q.E.D.

²¹In the case $\theta = 1$, there is k such that $g_1^k(x) \geq \varepsilon$ for each $x \geq \beta_\lambda \lambda^{-1}$.

C.4. Estimates on interim beliefs

Proposition C.2 below is the central step. For $k \in \{1, \dots, n-1\}$, we set $\psi_k(x) = x^k(1-x)^{n-k}$, and denote by \bar{g}_θ the function g_θ^β in the limit case $\beta = 0$. Proposition C.2 relates likelihood ratios to the average value of ψ_k and ψ_{n-k} along doubly infinite orbits of \bar{g}_1 .

PROPOSITION C.2: For each $k \in \{1, \dots, n-1\}$, one has

$$\limsup_{\lambda \rightarrow 0} \frac{p_k}{p_{n-k}} \leq \sup_{x \in (0,1)} \frac{\sum_{i \in \mathbf{Z}} \psi_k(\bar{g}_1^i(x))}{\sum_{i \in \mathbf{Z}} \psi_{n-k}(\bar{g}_1^i(x))}. \quad (\text{C.3})$$

We emphasize that in this statement, the interim belief p_k depends on the transition rate λ and the information acquisition strategy β_λ .

Exchanging p_k and p_{n-k} yields the inequality $\liminf_{\lambda \rightarrow 0} \frac{p_k}{p_{n-k}} \geq \inf_{x \in (0,1)} \frac{\sum_{i \in \mathbf{Z}} \psi_k(\bar{g}_1^i(x))}{\sum_{i \in \mathbf{Z}} \psi_{n-k}(\bar{g}_1^i(x))}$.

PROOF: Let $\varepsilon > 0$ and $k \in \{1, \dots, n-1\}$ be given. Given λ , we choose m (independent of λ) such that $g_1^{m+1}(\varepsilon) > 1 - \varepsilon$ and set $\varepsilon' = 1 - g_1^{m+1}(\varepsilon) \leq \varepsilon$. Note that

$$\frac{p_k}{p_{n-k}} = \frac{\int_0^1 \psi_k(x) dF_1(x)}{\int_0^1 \psi_{n-k}(x) dF_1(x)}. \quad (\text{C.4})$$

We write the numerator as

$$N_k := \int_0^\varepsilon \psi_k(x) dF_1(x) + \int_\varepsilon^{1-\varepsilon'} \psi_k(x) dF_1(x) + \int_{1-\varepsilon'}^1 \psi_k(x) dF_1(x).$$

Because of Lemma C.6 and since $\int_{1-\varepsilon'}^1 \psi_k(x) dF_1(x) = \int_0^{\varepsilon'} \psi_{n-k}(x) dF_0(x)$ by symmetry, there exists $C_0 < +\infty$ such that for every λ small, one has

$$N_k = c_{\lambda, \varepsilon} \lambda \varepsilon + \sum_{i=0}^m \int_{g_1^i(\varepsilon)}^{g_1^{i+1}(\varepsilon)} \psi_k(x) dF_1(x), \quad (\text{C.5})$$

for some $c_{\lambda, \varepsilon} \in [-C_0, C_0]$.

Let $i_0 = \max \{i : g^i(\varepsilon) \leq \frac{1}{2}\}$ and $z = g^{i_0}(\varepsilon)$.²² For each i , the change-of-variable formula for Stieltjes integrals yields

$$\int_{g_1^{i_0}(\varepsilon)}^{g_1^{i_0+1}(\varepsilon)} \psi_k(x) dF_1(x) = \int_z^{g_1(z)} \psi_k(g_1^{i-i_0}(x)) dF_1(g_1^{i-i_0}(x)).$$

Lemma C.2 and the continuous differentiability of ψ_k and of $g_1^{i-i_0}$ imply the existence of $C_1 < \infty$ such that

$$\int_z^{g_1(z)} \psi_k(g_1^{i-i_0}(x)) dF_1(g_1^{i-i_0}(x)) = \int_z^{g_1(z)} \psi_k(g_1^{i-i_0}(x)) dF_1(x) + d_{\lambda,\varepsilon} m k \lambda^2$$

for some $d_{\lambda,\varepsilon} \in [-C_1, C_1]$. Plugging into (C.5), we get

$$\frac{1}{\lambda} N_k = c_{\lambda,\varepsilon} \varepsilon + d_{\lambda,\varepsilon} m k \lambda + \sum_{i=-i_0}^{m-i_0} \frac{1}{\lambda} \int_z^{g_1(z)} \psi_k(g_1^i(x)) dF_1(x). \quad (\text{C.6})$$

For fixed ε , we deduce from (C.6) and its counterpart for N_{n-k} that

$$\frac{p_k}{p_{n-k}} = \frac{c_{\lambda,\varepsilon} \varepsilon + d_{\lambda,\varepsilon} m k \lambda + \frac{1}{\lambda} \int_z^{g_1(z)} \left[\sum_{i=-i_0}^{m-i_0} \psi_k(g_1^i(z)) \right] dF_1(x)}{c'_{\lambda,\varepsilon} \varepsilon + d'_{\lambda,\varepsilon} m k \lambda + \frac{1}{\lambda} \int_z^{g_1(z)} \left[\sum_{i=-i_0}^{m-i_0} \psi_{n-k}(g_1^i(z)) \right] dF_1(x)} \quad (\text{C.7})$$

for suitable constants $c'_{\lambda,\varepsilon}$ and $d'_{\lambda,\varepsilon}$.

We note that Lemma C.5 implies that for $a \in \{k, n-k\}$, the expression

$$\frac{1}{\lambda} \int_z^{g_1(z)} \psi_a(x) dF_1(x) \geq \left(\min_{x \in [\varepsilon, g_1(1-\varepsilon)]} \psi_a(x) \right) \times \min_{z \in [\varepsilon, g_1(\varepsilon)]} \frac{1}{\lambda} (F_1(g_1(z)) - F_1(x))$$

is bounded away from zero as $\lambda \rightarrow 0$.

²²Note that for λ small, i_0 does not depend on λ .

It thus follows from (C.7) that

$$\frac{p_k}{p_{n-k}} \leq \frac{\int_z^{g_1(z)} \left[\sum_{i=-i_0}^{m-i_0} \psi_k(g_1^i(z)) \right] dF_1(x)}{\int_z^{g_1(z)} \left[\sum_{i=-i_0}^{m-i_0} \psi_{n-k}(g_1^i(z)) \right] dF_1(x)} + C(\varepsilon + mk\lambda) \quad (\text{C.8})$$

for some constant C . We now note that

$$\frac{\int_z^{g_1(z)} \sum_{i=-i_0}^{m-i_0} \psi_k(g_1^i(x)) dF_1(x)}{\int_z^{g_1(z)} \sum_{i=-i_0}^{m-i_0} \psi_{n-k}(g_1^i(x)) dF_1(x)} \leq \sup_{x \in [z, g_1(z)]} \frac{\sum_{i=-i_0}^{m-i_0} \psi_k(g_1^i(x))}{\sum_{i=-i_0}^{m-i_0} \psi_{n-k}(g_1^i(x))}. \quad (\text{C.9})$$

For fixed ε (and hence fixed m and i_0), the right-hand side of (C.9) converges to

$$\sup_{x \in [z, \bar{g}_1(z)]} \frac{\sum_{i=-i_0}^{m-i_0} \psi_k(\bar{g}_1^i(x))}{\sum_{i=-i_0}^{m-i_0} \psi_{n-k}(\bar{g}_1^i(x))} \leq \sup_x \frac{\sum_{i \in \mathbf{Z}} \psi_k(\bar{g}_1^i(x))}{\sum_{i \in \mathbf{Z}} \psi_{n-k}(\bar{g}_1^i(x))},$$

as $\lambda \rightarrow 0$. Taking first the limit $\lambda \rightarrow 0$, then $\varepsilon \rightarrow 0$ in (C.8) yields the result. *Q.E.D.*

C.5. Conclusion

We have shown that given \hat{p} , there is λ_0 such that for $\lambda < \lambda_0$, there is a strategy β_λ for which $p_n = 1 - p_0 = \hat{p}$. To complete the proof, we need to show that provided \hat{p} is high enough, all interim beliefs p_k ($k = 1, \dots, n-1$) are such that $p_k \leq \hat{p}$ for λ small. This follows from Proposition C.2 and Lemma C.7 below.

LEMMA C.7: *One has $\sup_x \sum_{i \in \mathbf{Z}} \psi_k(\bar{g}_1^i(x)) < +\infty$ and $\inf_x \sum_{i \in \mathbf{Z}} \psi_k(\bar{g}_1^i(x)) > 0$.*

PROOF: For the lower bound, note that $\inf_x \sum_{i \in \mathbf{Z}} \psi_k(\bar{g}_1^i(x)) \geq \min_{x \in [\frac{1}{2}, \bar{g}_1(\frac{1}{2})]} \psi_k(x) > 0$.

For the upper bound, let $\varepsilon > 0$ be such that $a_1 := \min_{[0,\varepsilon]} \bar{g}'_1(x) > 1$ and $a_0 := \max_{[0,\varepsilon]} \bar{g}'_0(x) < 1$, and let m be s.t. $\bar{g}_1^m(\varepsilon) \geq 1 - \varepsilon$.

Then, for each $x \in (0, 1)$, one has

$$\begin{aligned} \sum_{i \in \mathbf{Z}} \psi_k(\bar{g}_1^i(x)) &\leq \sum_{i: \bar{g}_1^i(x) \leq \varepsilon} \psi_k(\bar{g}_1^i(x)) + \sum_{i: \bar{g}_1^i(x) \in (\varepsilon, 1-\varepsilon)} \psi_k(\bar{g}_1^i(x)) + \sum_{i: \bar{g}_1^i(x) \geq 1-\varepsilon} \psi_k(\bar{g}_1^i(x)) \\ &\leq \sum_{i: \bar{g}_1^i(x) \leq \varepsilon} \bar{g}_1^i(x) + m + \sum_{i: \bar{g}_1^i(x) \geq 1-\varepsilon} (1 - \bar{g}_1^i(x)) \\ &\leq \frac{a_1}{a_1 - 1} \varepsilon + k + \frac{1}{1 - a_0} (1 - \varepsilon) = C < \infty. \end{aligned}$$

Q.E.D.

APPENDIX D: CONTINUUM OF ACTIONS: PROPOSITION 4

Denote by $p_{a^{(1)}}$ and $p_{a^{(1)}a^{(2)}}$ the interim beliefs after sampling one or two actions. The action $a^{(1)}$ coincides with the posterior belief of the sampled agent, hence $\mathbf{E}[u(a^{(1)})]$ is an upper bound on the welfare and $\mathbf{E}[u(p_{a^{(1)}a^{(2)}})] \leq \mathbf{E}[u(a^{(1)})]$. On the other hand, taking transitions into account, one has

$$p_{a^{(1)}} = (1 - \lambda)a^{(1)} + \lambda(1 - a^{(1)}), \quad (\text{D.1})$$

and therefore, $\mathbf{E}[u(p_{a^{(1)}a^{(2)}})] \leq \mathbf{E}[u(p_{a^{(1)}})] + \lambda$.

Since u is quadratic, $u(p) = u(q) + (p - q)u'(q) + (p - q)^2$ for each p and q , so that

$$\mathbf{E}[u(p_{a^{(1)}a^{(2)}})] - \mathbf{E}[u(p_{a^{(1)}})] = \mathbf{E}[(p_{a^{(1)}a^{(2)}} - p_{a^{(1)}})(2p_{a^{(1)}} - 1)] + \mathbf{E}[(p_{a^{(1)}a^{(2)}} - p_{a^{(1)}})^2].$$

By iterated conditional expectations, the first expectation on the RHS is zero, so that $\mathbf{E}[(p_{a^{(1)}a^{(2)}} - p_{a^{(1)}})^2] \leq \lambda$ and similarly, $\mathbf{E}[(p_{a^{(1)}a^{(2)}} - p_{a^{(2)}})^2] \leq \lambda$. Using the inequality $(x - y)^2 \leq 2(x^2 + y^2)$, we get $\mathbf{E}[(p_{a^{(1)}} - p_{a^{(2)}})^2] \leq 2\lambda$. The result follows from (D.1).